

On the Canonical Height Gap for Polynomial Maps and Portraits of Preperiodic Points of Polynomials with Height 0

Haruki Imamura

Haruki Imamura, arXiv:2607.10160v1 (2026), 15 pages

<https://arxiv.org/abs/2607.10160v1>

One-sentence summary. The sparse homogeneous form of a polynomial map makes its resultant estimates uniform enough to bound the canonical–naïve height gap by $O(h(\varphi) + 1)$, independently of the degree up to bounded factors.

The problem and setup

Let $\varphi: \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be a polynomial map of degree $d \geq 2$ over a number field K . In homogeneous coordinates one may write

$$\varphi = (F(X_0, X_1) : X_1^d), \quad F = \sum_{j=0}^d a_j X_0^{d-j} X_1^j,$$

and $h(\varphi)$ is the projective height of the coefficients of this pair. For the logarithmic Weil height h on $\mathbf{P}^1$, the canonical height is

$$\widehat{h}_\varphi(P) = \lim_{n \rightarrow \infty} d^{-n} h(\varphi^n(P)).$$

The qualitative fact $\widehat{h}_\varphi = h + O(1)$ is classical. The issue is to control the constant explicitly without the linear-in- d loss present for a general rational map.

Main theorem (Imamura, Theorem 1.3). *For every $P \in \mathbf{P}^1(\overline{K})$,*

$$|\widehat{h}_\varphi(P) - h(P)| \leq \frac{d}{d-1} \left(\log 2 + \frac{\log d}{d} + h(\varphi) \right).$$

Since $d/(d-1) \leq 2$ and $(\log d)/(d-1)$ is bounded, this is genuinely $O(h(\varphi) + 1)$ uniformly in d . The polynomial hypothesis is essential to the argument: it is the monomial second coordinate X_1^d that creates the sparse matrices below.

Three proof ingredients

1. **Reduce the problem to a one-step error.** The standard height estimate gives

$$h(\varphi(P)) - dh(P) \leq \log(d+1) + h(\varphi).$$

Thus the missing input is a comparably sharp lower bound. Once both signs are controlled by a constant C , the definition of the canonical height turns the global comparison into a geometric-series calculation.

2. **Exploit the polynomial resultant.** Because F and X_1^d have no common projective zero, Bezout identities of degree $2d-1$ exist:

$$X_i^{2d-1} = g_{i0}F + g_{i1}X_1^d \quad (i = 0, 1),$$

where each g_{ij} has degree $d-1$. Cramer’s rule expresses their coefficients through minors of the Sylvester matrix of F and X_1^d . The paper’s Lemma 3.2 shows that the relevant staircase-shaped determinant is a sum of at most 2^{d-1} degree- d monomials. Consequently the joint projective height of the coefficients of all four g_{ij} is at most

$$2^{d-1} H(\varphi)^d.$$

This combinatorial estimate is the hardest point: a generic determinant count would lose the degree-uniform conclusion, whereas the monomial coordinate X_1^d forces precisely the needed sparsity.

3. **Pass from local inequalities to heights and telescope.** Applying the Bezout identities at every place v bounds $|P|_v^d$ by $|\varphi(P)|_v$ times the largest coefficient of the g_{ij} , with only the archimedean factor $2d$. Multiplication over all places gives

$$H(P)^d \leq 2^d d H(\varphi)^d H(\varphi(P)),$$

and hence

$$h(\varphi(P)) - dh(P) \geq -(d \log 2 + \log d + dh(\varphi)).$$

Together with the upper estimate this bounds the absolute one-step error by $C = d \log 2 + \log d + dh(\varphi)$. Finally,

$$\frac{h(\varphi^n(P))}{d^n} - h(P) = \sum_{i=1}^n \frac{h(\varphi^i(P)) - dh(\varphi^{i-1}(P))}{d^i},$$

so summing the geometric series and taking $n \rightarrow \infty$ proves the theorem.

Toy recurrence

If a real sequence satisfies $a_{n+1} = da_n + e_n$ with $|e_n| \leq C$, then $d^{-n}a_n - a_0 = \sum_{i=0}^{n-1} d^{-i-1}e_i$, and its limiting displacement is at most $C/(d-1)$. This is exactly the final height argument. The toy model removes all arithmetic geometry: the substantive work is proving a suitable C simultaneously at every place.

Height-zero application

If $\varphi \in \mathbf{Q}[X]$ has height zero, its coefficients lie in $\{-1, 0, 1\}$ and its leading coefficient is ± 1 . A rational preperiodic point α is therefore an integer (it is a rational algebraic integer), and $\widehat{h}_\varphi(\alpha) = 0$. The theorem gives $H(\alpha) \leq 8$ for $d = 2$ and $H(\alpha) < 5$ for $d \geq 3$.

It remains to turn these finite bounds into a universal list. Write $r = |\alpha|$. For a quadratic map, the triangle inequality gives $|\varphi(\alpha)| \geq r^2 - r - 1$; this is larger than r when $r \geq 4$, while $r = 3$ maps to absolute value at least 5 and then escapes. For $d \geq 3$,

$$|\varphi(\alpha)| \geq r^d - \sum_{i=0}^{d-1} r^i,$$

which is already at least $14 > r$ for $r \geq 3$. Reapplying the same estimate shows strict growth along the orbit. Thus every integer with $|\alpha| \geq 3$ escapes, proving Theorem 1.4:

$$\alpha \in \{0, \pm 1, \pm 2\}.$$

This is a universal necessary list, not a claim that all five points are preperiodic for each map. The paper combines it with the fact that integral cycles of a monic or antimonic polynomial have period at most two, then checks the remaining finite cases to classify and realize all possible portraits. The explicit realizations also make the range concrete: $X^3 + X$ has only 0 preperiodic among the five candidates; X^3 fixes $-1, 0, 1$; and for $X^2 - 1$ the points $0, -1$ form a 2-cycle while 1 maps into it.

Caveats and omissions

The note omits the paper's portrait diagrams and its case-by-case realization polynomials. It also does not claim sharpness of the height-gap constant. The coefficient height is the paper's fixed projective-coordinate height, so the statement should not be read as invariant under arbitrary conjugation of the dynamical system.