

Autoequivalences of Derived Categories of Moduli Spaces of Vector Bundles

Haotian Zuo

Haotian Zuo, arXiv:2605.28846v1 (2026), 7 pages

<https://arxiv.org/abs/2605.28846v1>

One-sentence summary. For fixed-determinant moduli of stable bundles on a curve of genus at least three, Fano reconstruction leaves no exotic derived symmetries: every autoequivalence is geometric, a line-bundle twist, or a shift.

Setup

Let C be a smooth projective curve of genus $g \geq 3$ over an algebraically closed field of characteristic zero. Fix $r \geq 2$ and a line bundle L of degree d with $\gcd(r, d) = 1$. Then

$$N = N_C(r, L)$$

is the smooth projective fine moduli space of stable rank- r bundles with determinant L . It is Fano, and $\text{Pic}(N) = \mathbf{Z}[\Theta]$ for the ample theta bundle Θ .

Define subgroups of $\text{Pic}(C) \times \text{Aut}(C)$ by

$$G_{[r]}^\circ = \{(\Lambda, \sigma) : \Lambda^r \simeq L \otimes \sigma^* L^{-1}\},$$

$$G_{[r]} = \{(\Lambda, \sigma) : \Lambda^r \simeq L \otimes \sigma^* L^{\pm 1}\}.$$

Main theorem (Zuo, Theorem 1.1 and Corollary 2.3). *The group of exact autoequivalences of $D^b(N)$ is*

$$\text{Aut } D^b(N) \simeq \begin{cases} \mathbf{Z}^2 \times G_{[r]}^\circ, & r \neq 2, \\ \mathbf{Z}^2 \times G_{[2]}, & r = 2. \end{cases}$$

The two infinite cyclic factors are generated by the shift $[1]$ and tensoring with Θ .

For $r \neq 2$, (Λ, σ) acts by $E \mapsto \sigma^* E \otimes \Lambda$. In rank two the larger group also contains the duality operation $E \mapsto \sigma^*(E^\vee) \otimes \Lambda$ whenever the determinant condition uses L^{+1} rather than L^{-1} .

Why the classification is so rigid

1. **Fano reconstruction reduces derived symmetry to ordinary geometry.** The Bondal–Orlov theorem applies because $-K_N$ is ample. Every exact autoequivalence is a composition of a shift, push-forward by an automorphism of N , and tensoring by a line bundle. Thus

$$\text{Aut } D^b(N) \simeq \mathbf{Z} \times (\text{Aut}(N) \ltimes \text{Pic}(N)).$$

This is the decisive input: unlike a Calabi–Yau or K3 category, no additional Fourier–Mukai symmetries can occur.

2. **The Picard action is trivial.** Ramanan’s computation gives $\text{Pic}(N) \simeq \mathbf{Z}[\Theta]$. If $a \in \text{Aut}(N)$, then $a^*\Theta$ is ample, so its class must be $[\Theta]$, not $-[\Theta]$. Hence the semidirect product is a direct product, and the Picard factor contributes the second copy of $\mathbf{Z}$.
3. **Automorphisms of the moduli space come from the curve.** The Kouvidakis–Pantev theorem identifies $\text{Aut}(N)$ with $G_{[r]}^\circ$ for $r \neq 2$ and with $G_{[2]}$ for $r = 2$. Pullback along an automorphism of C and tensoring by Λ must satisfy the displayed determinant equation. Rank two is special because dualization again has rank two and, after the compensating twist, preserves the fixed determinant. Combining this geometric classification with the first two steps proves the formula.

A second construction: extension correspondences

The paper also builds exact functors between moduli spaces of different ranks. After twisting, assume $0 < d < r$, and choose the unique lattice point (s, e) in the prescribed region satisfying

$$ds - re = 1, \quad t = r - s, \quad f = d - e.$$

For stable bundles $E \in M_C(s, e)$ and $F \in M_C(t, f)$, Lemma 3.1 proves that every non-split extension

$$0 \longrightarrow E \longrightarrow G \longrightarrow F \longrightarrow 0$$

is stable of rank r and degree d . The Bezout identity $ds - re = 1$ creates the narrow slope gap that excludes destabilizing subbundles in the three possible cases for their map to F .

On $M_C(s, e) \times M_C(t, f)$, the relative sheaf

$$\mathcal{E} = \mathcal{E}xt_{\pi_{12}}^1(\pi_{23}^* \mathcal{E}_2, \pi_{13}^* \mathcal{E}_1)$$

is locally free: stability gives $\text{Hom}(F, E) = 0$, Riemann–Roch makes $\dim \text{Ext}^1(F, E)$ constant, and Grauert’s theorem applies. The projective bundle $p : Z = \mathbf{P}(\mathcal{E}) \rightarrow M_C(s, e) \times M_C(t, f)$ parametrizes extension classes. Its universal extension has stable middle term, hence defines $q : Z \rightarrow M_C(r, d)$. The kernel $i_* \mathcal{O}_Z$ for $i = (p, q)$ yields the exact Fourier–Mukai type functor

$$\Phi \simeq Rq_* p^* : \mathbf{D}^b(M_C(s, e) \times M_C(t, f)) \longrightarrow \mathbf{D}^b(M_C(r, d)).$$

Restricting to compatible fixed determinants gives the analogous functor Ψ of Theorem 1.3.

Toy model and caveats

Projective space is the basic Fano model: its derived autoequivalences are generated by projective automorphisms, line-bundle twists, and shifts. The theorem says that $N_C(r, L)$ has the same rigidity pattern, while the finite or discrete factor $G_{[r]}$ remembers the curve and determinant.

The coprimality hypothesis makes the moduli spaces smooth and fine, and the autoequivalence formula stated here requires $g \geq 3$. For $g = 2$, rank two has a related formula without the extra dualization symmetry, while higher rank remains open in the paper. Finally, Theorems 1.2–1.3 assert that Φ and Ψ are exact; they do not claim these functors are fully faithful or equivalences.