

On the height boundedness of periodic and preperiodic points of dominant rational self-maps on projective varieties

Yohsuke Matsuzawa, Kaoru Sano

Yohsuke Matsuzawa and Kaoru Sano, arXiv:2603.09010v1 (2026), 30 pages

<https://arxiv.org/abs/2603.09010v1>

One-sentence summary. For a cohomologically hyperbolic rational self-map, periodic points whose entire orbits avoid a fixed proper closed set have uniformly bounded height.

The height question

Let X be a projective variety over $\overline{\mathbf{Q}}$, and let $f : X \dashrightarrow X$ be dominant. Write

$$X_f(\overline{\mathbf{Q}}) = \{P : f^n(P) \notin I_f \text{ for every } n \geq 0\}$$

for the points with well-defined forward orbit. If $d_i(f)$ is the i -th dynamical degree, then f is *cohomologically hyperbolic* when one $d_p(f)$ is strictly larger than every other dynamical degree. For the adjacent ratios

$$\mu_i = \frac{d_i(f)}{d_{i-1}(f)}, \quad \mu_{\dim X+1} = 0,$$

one then has $\mu_p > 1 > \mu_{p+1}$.

Main theorem (Matsuzawa–Sano, Theorem 1.8). *If f is cohomologically hyperbolic, there is a nonempty Zariski open subset $U \subset X$ such that*

$$\{P \in X_f(\overline{\mathbf{Q}}) : P \text{ is periodic and } O_f(P) \subset U\}$$

is height bounded.

Thus the conclusion is uniform even over algebraic points of unbounded degree. The qualification $O_f(P) \subset U$ is essential to the proof: the height inequalities come from a big divisor and hold only away from its exceptional locus.

The divisor inequality

Fix an ample divisor L and a corresponding height $h = h_L$. Choose $\varepsilon < 1$ close to 1 and a large m so that

$$\alpha = (\varepsilon^2 \mu_p)^m > 1, \quad \beta = (\varepsilon^{-2} \mu_{p+1})^m < 1.$$

A pullback positivity theorem gives the big class

$$(f^{2m})^*L + \alpha\beta L - (\alpha + \beta)(f^m)^*L$$

on a common birational resolution of f^m and f^{2m} . A height attached to a big divisor is bounded below on some nonempty open set. After shrinking that set to U , one obtains

$$h(f^{2m}Q) + \alpha\beta h(Q) - (\alpha + \beta)h(f^mQ) \geq c \quad (Q \in U(\overline{\mathbf{Q}})).$$

This is the geometric heart of the proof: cohomological hyperbolicity separates the two characteristic roots of a second-order height recurrence, placing one above and one below 1.

How periodicity kills both modes

Put

$$c_1 = \frac{c}{1-\alpha}, \quad c_2 = \frac{c}{1-\beta},$$
$$\Phi = h \circ f^m - \beta h - c_1, \quad \Psi = h \circ f^m - \alpha h - c_2.$$

The divisor inequality factors into the two recursive estimates

$$\Phi(f^m Q) \geq \alpha \Phi(Q), \quad \Psi(f^m Q) \geq \beta \Psi(Q).$$

Let P be periodic with $O_f(P) \subset U$. The values of Φ on its orbit form a finite set. Since $\alpha > 1$, the first inequality would make any positive value grow indefinitely, so

$$\Phi(P) \leq 0, \quad h(f^m P) \leq \beta h(P) + c_1.$$

Choose $N \geq 1$ with $f^{mN}(P) = P$. Iterating the second inequality gives $\Psi(P) \geq \beta^N \Psi(P)$. Because $0 < \beta < 1$, this forces

$$\Psi(P) \geq 0, \quad h(f^m P) \geq \alpha h(P) + c_2.$$

Combining the two bounds eliminates $h(f^m P)$:

$$h(P) \leq \frac{c_1 - c_2}{\alpha - \beta}.$$

The right side depends on f, L , and U , but not on the periodic point.

Toy recurrence

For a periodic numerical sequence, an inequality governed by $(T - \alpha)(T - \beta)$, with $\alpha > 1 > \beta > 0$, cannot retain an expanding component: periodicity prevents its growth. Running once around the cycle also prevents a negative contracting component. The functions Φ and Ψ implement exactly this factorization for heights; the big-divisor theorem is what supplies the recurrence outside the exceptional set.

Scope and counterpoints

The paper first disproves a broader affine expectation. The automorphism

$$(x, y, z) \mapsto (y, x + y - y^3, 2z - y)$$

of $\mathbf{A}^3$ has finitely many n -periodic points for every n , yet a Zariski-dense sequence of periodic points has height tending to infinity (Theorems 1.3 and 2.1). Here $d_1 = d_2 = 3$, so cohomological hyperbolicity fails.

The theorem does not bound all periodic points without the orbit condition. It also does not extend formally to preperiodic points: Theorem 1.9 constructs a cohomologically hyperbolic map with a Zariski-dense backward chain whose heights diverge. The authors carefully note that this is not yet a counterexample to the strongest possible open-set statement for preperiodic points, and finiteness over a fixed number field remains open.