

On Fano indices of weighted projective spaces

Haidong Liu

Haidong Liu, arXiv:2608.03434v1 (2026), 15 pages

<https://arxiv.org/abs/2608.03434v1>

One-sentence summary. Canonical singularities impose a sharp Sylvester-sequence upper bound on the Fano index of every well-formed weighted projective space.

From weights to the index

Let

$$X = \mathbf{P}(a_1, \dots, a_{n+1}), \quad a_1 \geq \dots \geq a_{n+1} > 0,$$

be well formed and have canonical singularities. Since $\mathcal{O}_X(1)$ generates the Weil divisor class group,

$$-K_X \sim_{\mathbf{Q}} (a_1 + \dots + a_{n+1})\mathcal{O}_X(1), \quad q(X) = a_1 + \dots + a_{n+1}.$$

Define the Sylvester sequence and its shifted form by

$$s_1 = 2, \quad s_i = s_1 \cdots s_{i-1} + 1, \quad y_i = s_i - 1.$$

Thus $s_1, s_2, s_3, s_4 = 2, 3, 7, 43$, and

$$\frac{1}{s_1} + \dots + \frac{1}{s_{i-1}} = 1 - \frac{1}{y_i}.$$

Main theorem (Liu, Theorem 1.2). *The Fano index of X satisfies*

$$q(X) \leq y_n(2y_n - 1) = (s_n - 1)(2s_n - 3).$$

The bound is sharp. It also applies to any n -dimensional $\mathbf{Q}$ -factorial toric Fano variety of Picard number one with canonical singularities (Corollary 1.3): such a variety has a finite cover, étale in codimension one, by a well-formed weighted projective space, and pullback cannot decrease the relevant divisibility bound.

Two arithmetic encodings

Set $q = \sum a_i$ and $\beta_i = a_i/q$. The vector $(\beta_1, \dots, \beta_{n+1})$ is the barycentric-coordinate vector of the unique interior lattice point of an integral simplex attached to X ; its normalized volume is q . The simplex supplies product-sum inequalities

$$\beta_1 \cdots \beta_j \leq \beta_{j+1} + \dots + \beta_{n+1}$$

and sharp lower or upper bounds for selected products.

The singularities are encoded by the integer-valued age function

$$\text{age}(k) = \sum_{i=1}^{n+1} \left\{ \frac{a_i k}{q} \right\} = k - \sum_{i=1}^{n+1} \left\lfloor \frac{a_i k}{q} \right\rfloor.$$

Well-formedness gives $1 \leq \text{age}(k) \leq n$ for $0 < k < q$. Kasprzyk's criterion says that canonical singularities are equivalent to

$$\text{age}(k) \neq n \quad (2 \leq k \leq q-2).$$

The proof manufactures exactly such a forbidden age if q is too large.

The contradiction in three steps

Assume $q > y_n(2y_n - 1)$.

1. **The two smallest weights become rigid.** Simplex product bounds and a refined product–sum inequality first show

$$a_n = a_{n+1} = y_n$$

(Lemmas 3.1–3.2). Any larger a_n , or any smaller a_{n+1} , makes the lower product bound incompatible with $\beta_1 \cdots \beta_n \leq 1/q$.

2. **Majorization controls every remaining weight.** For $1 \leq i \leq n - 1$, Lemma 3.3 proves

$$\frac{a_i}{q} < \frac{2y_n}{s_i(2y_n - 1)}.$$

The difficult cases compare $(\beta_1, \dots, \beta_{n-1})$ with a small perturbation of $(1/s_1, \dots, 1/s_{n-1})$. The comparison vector majorizes the barycentric vector, so Karamata’s inequality for the concave function $\log$ gives the needed product lower bound. If one β_i crossed the displayed threshold, that lower bound would contradict $y_n \beta_1 \cdots \beta_{n-1} \leq 1$.

3. **A single age value detects the contradiction.** Take $k = 2y_n - 1$. The last two weights contribute zero to the floor sum, while Lemma 3.3 and the integrality of $2y_n/s_i$ give

$$\left\lfloor \frac{ka_i}{q} \right\rfloor \leq \frac{2y_n}{s_i} - 1 \quad (1 \leq i \leq n - 1).$$

Using the Sylvester identity,

$$\begin{aligned} \text{age}(2y_n - 1) &\geq 2y_n - 1 - \sum_{i=1}^{n-1} \left(\frac{2y_n}{s_i} - 1 \right) \\ &= n. \end{aligned}$$

Since age is at most n , equality holds, contradicting the canonical singularity criterion. Hence the assumed strict inequality for q is impossible.

Dimension four and scope

For $n = 4$, $y_4 = 42$, so the theorem reads

$$q(X) \leq 42 \cdot 83 = 3486.$$

The proof says that a violation would force $\text{age}(83) = 4$, a forbidden cyclic-quotient signature. This numerical case cleanly displays the role of the Sylvester sequence.

The result concerns well-formed weighted projective spaces and, through the finite-cover argument, $\mathbf{Q}$ -factorial toric Fano varieties of Picard number one. The corresponding bound for arbitrary canonical Fano varieties remains the paper’s Conjecture 1.1. The separate computational classification of four-dimensional index values (Theorem 1.4) and the conjectural comparison with terminal Calabi–Yau indices are not used in the proof summarized here.