

# On products of K-moduli spaces

Theodoros S. Papazachariou

Theodoros S. Papazachariou, arXiv:2403.04544v2 (2024), 25 pages

<http://arxiv.org/abs/2403.04544v2>

**One-sentence summary.** When two K-semistable Fano varieties share no simple factor, taking products identifies the corresponding reduced K-moduli components, both as stacks and as good moduli spaces.

## The product question

A  $\mathbf{Q}$ -Fano variety is a normal projective variety with klt singularities and ample  $\mathbf{Q}$ -Cartier anticanonical divisor. K-semistable  $\mathbf{Q}$ -Fano varieties of fixed dimension and volume form an Artin stack  $\mathcal{M}_{n,V}^K$ , whose good moduli space parametrizes K-polystable objects.

For a K-semistable Fano variety  $X_i$ , the paper writes  $\mathcal{M}_{X_i}^K$  for the *reduced connected component* containing the deformations and K-semistable degenerations of  $X_i$ , and  $M_{X_i}^K$  for its good moduli space. If  $X = X_1 \times X_2$ , fiber product of families defines

$$\text{Prod} : \mathcal{M}_{X_1}^K \times \mathcal{M}_{X_2}^K \longrightarrow \mathcal{M}_X^K, \quad ([Y], [Z]) \longmapsto [Y \times Z].$$

Call a  $\mathbf{Q}$ -Fano variety *simple* if it is not a product of lower-dimensional  $\mathbf{Q}$ -Fano varieties.

**Main theorem** (Papazachariou, Theorem 5.3). *Let  $X_1 \not\cong X_2$  be K-semistable Fano varieties such that no simple factor in the product decomposition of  $X_1$  is isomorphic to a simple factor of  $X_2$ . Then*

$$\mathcal{M}_{X_1}^K \times \mathcal{M}_{X_2}^K \xrightarrow{\sim} \mathcal{M}_{X_1 \times X_2}^K, \quad M_{X_1}^K \times M_{X_2}^K \xrightarrow{\sim} M_{X_1 \times X_2}^K.$$

This upgrades the pointwise fact that a product is K-semistable exactly when its factors are: it identifies whole moduli components and their families.

## Rigidity of the factors

Theorem 2.11 proves that every  $\mathbf{Q}$ -Fano variety has a unique decomposition into simple  $\mathbf{Q}$ -Fano factors, up to permutation and factorwise isomorphism. Kawamata–Viehweg vanishing and Künneth give

$$H^i(X, \mathcal{O}_X) = 0 \ (i > 0), \quad \text{Pic}(X_1 \times \cdots \times X_m) \cong \prod_i \text{Pic}(X_i).$$

Pulling back an ample bundle from a putative factor therefore selects a subproduct through its semiample linear system. Simplicity forces that subproduct to be a single factor. This is what prevents a product point from having a second, incompatible factorization.

## Why deformations also split

The local geometric input is that a flat family with central fiber  $X_1 \times X_2$  splits analytically, and then étale-locally, as a fiber product (Lemma 2.7 and Theorem 3.1). The stronger deformation-theoretic statement is Theorem 3.4:

$$\mathrm{Def}_{X_1}^{\mathrm{qG}} \times \mathrm{Def}_{X_2}^{\mathrm{qG}} \xrightarrow{\sim} \mathrm{Def}_{X_1 \times X_2}^{\mathrm{qG}}.$$

For the canonical covering stacks, the cotangent complex splits as

$$L_{X_1 \times X_2} \simeq p_1^* L_{X_1} \oplus p_2^* L_{X_2}.$$

Vanishing of higher cohomology of the structure sheaves identifies every relevant Ext group with the direct sum of the groups for the factors. The tangent spaces and obstruction theory therefore split; induction over small derived Artin algebras shows that every  $\mathbb{Q}$ -Gorenstein deformation of the product exists uniquely as a product deformation. Consequently the product locus is open, and a connected reduced component containing one product consists entirely of products.

## Globalizing the local picture

Three further steps turn this local statement into the theorem.

1. Products preserve klt singularities, K-semistability, and Kollár’s base-change condition, so fiber product really defines a morphism of K-moduli stacks (Proposition 4.2). Products of the relevant quotient stacks also have the product of their good moduli spaces (Proposition 5.2).
2. Unique simple-factor decomposition and the hypothesis excluding common factors make  $\mathrm{Prod}$  injective on objects and on stabilizers. Together with deformation splitting, this gives a representable, separated open immersion that sends closed points to closed points.
3. On good moduli spaces, properness plus quasi-finiteness makes the descended map finite. The stack morphism is then finite as well; a finite open immersion is an isomorphism, by the stack version of Zariski’s main theorem.

## Repeated factors and scope

The hypothesis records a genuine symmetry. For distinct rigid factors such as  $\mathbf{P}^1$  and  $\mathbf{P}^2$ , the product component has no ambiguity from permuting factors. For a repeated simple factor  $X_1 = X_2$ , exchanging the two copies survives: Theorem 5.4 describes the product map as an  $S_2$ -gerbe and proves that it is étale, rather than an isomorphism from the ordered product.

The theorem is over  $\mathbf{C}$  and concerns specified reduced connected components, not the entire K-moduli stack. Nonreduced local structure, components with repeated or shared simple factors, and the paper’s separate log-Fano and wall-crossing applications require the additional statements developed later in the paper.