

Arithmetic degree and its application to Zariski dense orbit conjecture

Yohsuke Matsuzawa, Junyi Xie

Yohsuke Matsuzawa and Junyi Xie, arXiv:2409.06160v3 (2024), 20 pages

<http://arxiv.org/abs/2409.06160v3>

One-sentence summary. Points whose arithmetic degree is arbitrarily close to the first dynamical degree are adelically dense; for a birational map with $\lambda_1 > \lambda_3$, this forces a Zariski-dense orbit whenever no nonconstant rational function is invariant.

Height growth versus geometric growth

Let $f: X \dashrightarrow X$ be a dominant rational self-map of a quasi-projective variety over $\overline{\mathbf{Q}}$. Write X_f for the points whose forward orbit avoids every indeterminacy locus. After choosing a projective closure and an ample Weil height h_L , set

$$\alpha_f(x) = \lim_{n \rightarrow \infty} \max\{1, h_L(f^n(x))\}^{1/n}, \quad \lambda_i(f) = \lim_{n \rightarrow \infty} ((f^n)^* L^i \cdot L^{\dim X - i})^{1/n}.$$

The first limit need not be known to exist for every rational map and point; the second does exist and is birationally invariant.

Main theorem (Matsuzawa–Xie, Theorems 3.1 and 4.1). *For every $\varepsilon > 0$, the set of $x \in X_f(\overline{\mathbf{Q}})$ for which $\alpha_f(x)$ exists and*

$$\alpha_f(x) \geq \lambda_1(f) - \varepsilon$$

is dense for the adelic topology, hence Zariski dense.

If X is projective, f is birational, and $\lambda_3(f) < \lambda_1(f)$ (vacuously when $\dim X \leq 2$), then the Zariski dense orbit conjecture holds for f . More precisely, if $\mathbf{k}(X)^f = \mathbf{k}$, points with Zariski-dense orbit are adelically dense.

In dimension three, $\lambda_3(f) = 1$ for a birational map, so $\lambda_1(f) > 1$ suffices. This replaces a previously used full cohomological-hyperbolicity hypothesis by a comparison of just two dynamical degrees.

The height recursion

Put $\mu_i = \lambda_i/\lambda_{i-1}$; log concavity gives $\mu_1 \geq \cdots \geq \mu_{\dim X}$. Choose the last p in the initial plateau $\mu_1 = \cdots = \mu_p > \mu_{p+1}$ and $\zeta < 1$ close to 1. After replacing f by a suitable iterate, a bigness estimate for pullbacks produces, on a dense open set, a two-step inequality of the form

$$h_L(f^{2m}x) - \zeta^{-2m} \mu_{p+1}^m h_L(f^m x) - c \geq \zeta^{2m} \mu_p^m (h_L(f^m x) - \zeta^{-2m} \mu_{p+1}^m h_L(x) - c).$$

Once the parenthesized initial quantity is positive, iteration gives exponential height growth at rate at least $\zeta^2 \mu_p = \zeta^2 \lambda_1(f)$.

How the initial positive value is found

The difficult point is not the recurrence but its seed. Three ingredients work together.

1. **Adelic control of the orbit.** Inside any prescribed nonempty adelic open set, earlier dynamical Mordell–Lang results provide a smaller adelic open set of points whose orbits are infinite, remain in the good locus, and satisfy DML. The last condition later guarantees existence of the arithmetic-degree limit.
2. **A curve of maximal degree growth.** For $g = f^m$, successive hyperplane sections through a good point give an irreducible curve C avoiding I_{g^ℓ} and satisfying

$$\deg((g^\ell|_C)^*L) = ((g^\ell)^*L \cdot L^{\dim X-1}).$$

The right side grows nearly as $\mu_p^{m^\ell}$, so comparison of heights on the normalization of C forces at least one large multiplicative jump $h_L(g^{i+1}x)/h_L(g^i x)$.

3. **Unbounded height in every adelic open set.** Proposition 2.3 proves this using bounded-degree algebraic points and Northcott finiteness. It lets one choose $x \in C$ high enough that the error terms cannot erase that jump. The recurrence then yields $\alpha_f(x) \geq \zeta^2 \lambda_1(f) \geq \lambda_1(f) - \varepsilon$.

For comparison, on $\mathbf{P}^1$ a morphism of degree d satisfies $h(f^n x) = d^n h(x) + O(d^{n-1})$; a positive starting height immediately reveals the rate d . The curve and adelic arguments manufacture an analogue of that favorable starting point despite indeterminacy and several cohomological growth rates.

From fast height growth to a dense orbit

Assume now that f is birational and has no invariant nonconstant rational function. There are only finitely many totally invariant hypersurfaces, so choose x outside them with $\alpha_f(x) > \lambda_3(f)$. If its orbit closure were proper, each periodic irreducible component W would have codimension at least two. For a suitable iterate,

$$\alpha_f(x)^m \leq \lambda_1(f^m|_W) \leq \lambda_{1+\text{codim } W}(f^m) \leq \lambda_3(f)^m,$$

where the last step is log concavity together with $\lambda_1(f) > \lambda_3(f)$. This contradicts the choice of x .

Scope

The approximation theorem retains an arbitrary ε ; the paper explicitly asks whether equality $\alpha_f(x) = \lambda_1(f)$ always occurs. The dense-orbit application requires birationality and the absence of an invariant rational fibration. Its conclusion is existence (indeed adelic density) of suitable algebraic points, not a classification of all orbits.