

Quantitative Equidistribution of Small Points for Canonical Heights

Jit Wu Yap

Jit Wu Yap, arXiv:2410.21679v1 (2024), 49 pages

<http://arxiv.org/abs/2410.21679v1>

One-sentence summary. Outside an explicitly bounded exceptional hypersurface, the discrepancy between a Galois orbit and the canonical measure is controlled by canonical height; the estimate yields exponential equidistribution of periodic points for polarized endomorphisms of surfaces.

Making Yuan's theorem quantitative

Let X be a smooth projective variety over a number field K , and let $\varphi: X \rightarrow X$ be polarized: for an ample line bundle L , $\varphi^*L \simeq L^{\otimes d}$ with $d \geq 2$. Write $\widehat{h}_\varphi$ for the associated canonical height. At an archimedean place v , the canonical metric determines an equilibrium measure $\mu_{\varphi,v}$ on $X(\mathbf{C}_v)$. If F_x denotes the $\text{Gal}(\overline{K}/K)$ -orbit of x , Yuan's qualitative theorem says that generic sequences with $\widehat{h}_\varphi(x) \rightarrow 0$ equidistribute to $\mu_{\varphi,v}$.

Main theorem (Yap, Theorem 1.2). *There are constants $c_1 = c_1(K, \varphi)$, $c_2 = c_2(\varphi)$, and $c_3 = c_3(K, \varphi)$ such that, for every $\varepsilon > 0$ and every smooth $f: X(\mathbf{C}_v) \rightarrow \mathbf{R}$, there is a hypersurface $H(f, \varepsilon)$ over K satisfying*

$$\deg_L H(f, \varepsilon) \leq c_1 \varepsilon^{-c_2}.$$

For every $x \in X(\overline{K}) \setminus H(f, \varepsilon)$,

$$\left| \frac{1}{|F_x|} \sum_{y \in F_x} f(y) - \int_X f d\mu_{\varphi,v} \right| \leq c_{f,3} \left(\frac{\widehat{h}_\varphi(x)}{\varepsilon} + c_3 \varepsilon \right),$$

where $c_{f,3}$ is controlled by derivatives of f through order three.

The degree bound quantifies the theorem's genericity requirement. To make the discrepancy $O(\varepsilon)$ it is enough to have $\widehat{h}_\varphi(x) = O(\varepsilon^2)$ and to avoid a hypersurface whose degree grows only polynomially in ε^{-1} . For an abelian variety with its Néron–Tate height, the canonical metric is smooth and the paper obtains the uniform exponent $c_2 = 8$.

Regularizing the canonical metric

Embed X into projective space so that φ extends to a projective endomorphism. Its homogeneous Green function supplies a semipositive adelic metric on L , but at an archimedean place this metric is generally only Hölder continuous. For $\varepsilon = d^{-k}$ and a large tensor power $L_n = L^{d^n}$, the proof regularizes the local metric potentials at a scale depending on ε . The resulting smooth metric h' obeys schematically

$$\|h' - h\|_\infty = O_\varphi(d^n \varepsilon^2), \quad dd^c h' \geq d^n \varepsilon^2 \omega_{\text{FS}},$$

while its derivatives grow by only a fixed power of ε^{-1} . The small Fubini–Study correction in the second inequality is what permits analytic estimates for positive line bundles.

Volumes produce the exceptional divisor

The proof now perturbs the metric at v by the test function f , through the factor $\exp(-d^n \varepsilon f)$. Arithmetic Hilbert–Samuel theory controls the covolume of integral sections. Distortion-function estimates compare the perturbed and unperturbed L^2 unit balls, and Gromov’s inequality transfers the result to supremum norms. After replacing the regularized Monge–Ampère measure by $\mu_{\varphi,v}$, the central volume estimate has leading term

$$\frac{1}{d^n \dim H^0(X, L_n)} \log \frac{\text{vol } B_n(f)}{\text{covol } M_n} \gtrsim \varepsilon \int_X f d\mu_{\varphi,v} - O_{\varphi}(\varepsilon^2).$$

Normalize f to have integral zero and bounded C^3 size. Taking $n = O_{\varphi}(k)$ makes the normalized volume positive enough for Minkowski’s theorem, which yields a nonzero global section s that is small at every place for the perturbed adelic metric. For $x \notin \text{div}(s)$, positivity of the height computed with s gives

$$\frac{1}{|F_x|} \sum_{y \in F_x} f(y) \geq -\frac{\widehat{h}_{\varphi}(x)}{\varepsilon} - c_3 \varepsilon.$$

Applying the argument to $-f$ gives the other inequality. Finally, $H(f, \varepsilon) = \text{div}(s) \cap X$ has degree d^n ; because $n = O(k)$ and $\varepsilon = d^{-k}$, this is at most a fixed power of ε^{-1} .

Dynamical consequences and scope

On a smooth projective surface, the exceptional curve contains relatively few points of period n : Bézout bounds handle nonperiodic components, while normalization and Riemann–Hurwitz control periodic ones. Combining these counts with the theorem gives constants $C > 0$ and $\lambda > 1$ such that the period- n points satisfy

$$\left| \frac{1}{|\text{Per}_n|} \sum_{x \in \text{Per}_n} f(x) - \int f d\mu_{\varphi,v} \right| \leq C c_{f,3} \lambda^{-n}.$$

The paper also derives exponential lower bounds for fields containing periodic points and polynomial-degree hypersurfaces containing preperiodic points of bounded algebraic degree. The main estimate is archimedean, requires smooth X and smooth test functions, and allows the exceptional hypersurface to depend on both f and ε .