

Global smoothing of singular Fano and Calabi-Yau varieties

Anda Tenie

Anda Tenie, arXiv:2602.07615v2 (2026), 25 pages

<http://arxiv.org/abs/2602.07615v2>

One-sentence summary. Du Bois–Nakano vanishing globalizes local deformation directions, showing that singular Fano varieties with isolated Du Bois lci singularities can be deformed until only 1-rational singularities remain—and can be smoothed when all initial singularities are 1-irrational.

The smoothing problem

Here a singular Fano variety is a projective complex variety Y of dimension $n \geq 3$ with isolated Du Bois local-complete-intersection (lci) singularities and ample ω_Y^{-1} . It is *smoothable* if it is the special fibre of a flat analytic family over a disk whose general fibre is smooth. Local smoothability alone does not solve the problem: first-order local directions need not come from one global deformation.

For a singular germ (X, x) , take a strong log resolution $\pi: \widehat{X} \rightarrow X$ with exceptional divisor E . Under the paper’s hypotheses, the singularity is 1-rational precisely when

$$H^{n-2}(\widehat{X}, \Omega_{\widehat{X}}^1(\log E)) = 0.$$

Equivalently, by duality, $H_E^2(\Omega_{\widehat{X}}^{n-1}(\log E)(-E)) = 0$; a failure of this vanishing is called 1-irrational.

Main theorem (Tenie, Theorems 3.1 and 3.7). *Let Y be a singular Fano variety with isolated Du Bois lci singularities. Then $T_Y^i = 0$ for $i \geq 2$, so its deformations are unobstructed. Moreover, Y admits a deformation to a Fano variety having only 1-rational singularities. If none of the singularities of Y is 1-rational, then Y is smoothable, and every small smoothing is Fano.*

Thus the first conclusion improves the local type even when some 1-rational points persist. The stronger hypothesis in the second conclusion says that every singular point is among those the construction can remove.

Why deformations are unobstructed

For an lci variety, $T_Y^i = \text{Ext}^i(\Omega_Y^1, \mathcal{O}_Y)$. Serre duality reduces its vanishing for $i \geq 2$ to a statement about $H^{n-i}(\Omega_Y^1 \otimes \omega_Y)$, where Ω_Y^1 is the first graded piece of the Du Bois complex. A hypercohomology spectral sequence and Steenbrink’s vanishing for isolated lci singularities inject this group into the corresponding Du Bois hypercohomology. Since ω_Y^{-1} is ample, a dual Nakano-type vanishing theorem kills the latter group. Hence every first-order global direction can be integrated to an actual small deformation.

From global classes to simultaneous local smoothing

Let $Z = \text{Sing}(Y)$, $V = Y \setminus Z$, and $\pi: \widehat{Y} \rightarrow Y$ be a strong log resolution with exceptional divisor E . Schlessinger's identifications compare global and local first-order deformations:

$$T_Y^1 \simeq H^1(V, T_V), \quad H^0(\mathcal{T}_Y^1) \simeq \bigoplus_{x \in Z} H^1(U_x, T_{U_x}).$$

The restriction map between these spaces is usually not surjective, so the proof inserts local cohomology. Set

$$\mathcal{G} = \Omega_{\widehat{Y}}^{n-1}(\log E)(-E) \otimes \pi^* \omega_Y^{-1}.$$

Local cohomology sequences and excision give a commutative diagram linking T_Y^1 to

$$H_E^2(\mathcal{G}) \simeq \bigoplus_{x \in Z} H_{E_x}^2 \left(\Omega_{\widehat{X}_x}^{n-1}(\log E_x)(-E_x) \right).$$

The crucial global group vanishes:

$$H^2(\widehat{Y}, \mathcal{G}) = \mathbb{H}^2(\Omega_Y^{n-1} \otimes \omega_Y^{-1}) = 0,$$

again by Du Bois–Nakano vanishing. Exactness therefore makes the map from global deformations onto this direct sum of local defect spaces surjective.

At a 1-irrational point the corresponding summand is nonzero. A local criterion (Lemma 2.10) says that a first-order class whose image is not trapped in the maximal-ideal multiple of the local deformation module smooths that point. Surjectivity lets one choose a single global direction with this property simultaneously at every 1-irrational point. The unobstructedness theorem integrates it. Semicontinuity of the minimal exponent preserves the already 1-rational points, so the new fibre has only 1-rational singularities; if there were none initially, all singular points are removed. Openness of ampleness keeps the nearby fibres Fano.

Calabi–Yau counterpart and limits

The same paper proves parallel results for singular Calabi–Yau varieties. Without an extra global condition one can deform to 1-Du Bois singularities; with $h^{n-1,2}(Y) = h^{1,n-2}(Y)$ one can deform to 1-rational singularities and, when all points are 1-irrational, smooth completely. For threefolds this Hodge–Du Bois equality recovers the role of familiar factoriality conditions. The results require isolated Du Bois lci singularities and dimension at least three; the theorem does not claim that arbitrary singular Fano or Calabi–Yau varieties are smoothable.