

Chern classes of Deligne-Mumford stacks and their coarse moduli spaces

Hsian-Hua Tseng

arXiv:0709.0034v2, 8 pages

<http://arxiv.org/abs/0709.0034v2>

One-sentence summary. Tseng identifies the Chern–Schwartz–MacPherson class of a quotient-singular coarse moduli space with tangent Chern data on the inertia stack, and its stringy Chern class with the corresponding data on the double inertia stack.

Two viewpoints on a singular coarse space

Let $\mathcal{X}$ be a smooth separated Deligne–Mumford stack of finite type over $\mathbf{C}$, with projective coarse moduli variety X and structure map $\pi : \mathcal{X} \rightarrow X$ (Assumption 2.1). The variety X has quotient singularities. Its Chern–Schwartz–MacPherson class $c^{SM}(X)$ is defined intrinsically from the singular variety, whereas $\mathcal{X}$ has an honest tangent bundle.

The inertia stack

$$I\mathcal{X} = \mathcal{X} \times_{\mathcal{X} \times \mathcal{X}} \mathcal{X}$$

records a point together with an automorphism. Write $p : I\mathcal{X} \rightarrow \mathcal{X}$ for the projection. Its own inertia stack $II\mathcal{X} := I(I\mathcal{X})$ records two commuting automorphisms; let $q : II\mathcal{X} \rightarrow \mathcal{X}$ be the natural composite projection.

Main theorem (Tseng, Theorems 3.4 and 4.1). *Under Assumption 2.1,*

$$\pi_* p_* c(T_{I\mathcal{X}}) = c^{SM}(X).$$

If IX is the coarse moduli space of $I\mathcal{X}$ and $\bar{p} : IX \rightarrow X$, then the stringy constructible function satisfies $\Phi_X = \bar{p}_ \mathbf{1}_{IX}$, and*

$$c_{\text{str}}(X) = \pi_* q_* c(T_{II\mathcal{X}}).$$

All classes are understood in rational Chow groups.

Thus singular characteristic classes on X are recovered from ordinary tangent bundles on smooth stacks carrying stabilizer data. The appearance of double inertia in the stringy formula is structural, not decorative.

The quotient model and the key function identity

For a local quotient $\mathcal{X} = [M/G]$ by a finite group,

$$I[M/G] = \coprod_{(g)} [M^g / C_G(g)],$$

where (g) runs over conjugacy classes. Tseng uses the stack pushforward of constructible functions, whose isotropy weights differ from ordinary fiber counting. The central calculation is Proposition 3.2:

$$\pi_* p_* \mathbf{1}_{I\mathcal{X}} = \mathbf{1}_X.$$

Locally it is the orbit–stabilizer cancellation encoded by

$$\frac{1}{|G|} \sum_{g \in G} \mathbf{1}_{M^g} = \sum_{(g)} \frac{\mathbf{1}_{M^g}}{|C_G(g)|}.$$

The factor $|C_G(g)|$ introduced when M^g is pushed to $[M^g / C_G(g)]$ cancels the denominator. An étale quotient chart then globalizes the identity. In the toy case $\mathcal{X} = BG$, the equality $\sum_{(g)} |C_G(g)|^{-1} = 1$ is the whole mechanism in miniature.

Four proof steps

1. **Compatible pushforwards.** Lemma 2.3 computes the pushforward of a characteristic function under a finite proper representable étale map. Comparing this definition with the equivariant one on quotient presentations ensures that functoriality can be used without losing stabilizer factors.
2. **From functions to Chern classes.** Apply the natural transformation C_* from constructible functions on quotient stacks to homology/Chow classes to Proposition 3.2. Naturality commutes with p_* and π_* , while normalization on the smooth stack $I\mathcal{X} = [W/H]$ gives $C_*(\mathbf{1}_{I\mathcal{X}}) = c(T_{I\mathcal{X}})$. This proves the CSM formula.
3. **Identify the stringy function.** The identity $\Phi_X = \bar{p}_*\mathbf{1}_{IX}$ is checked étale-locally. Relative canonical divisors commute with étale base change, reducing to a finite quotient $X = M/G$, where the required formula is the established quotient calculation used by the paper.
4. **Iterate inertia.** By definition the inertia of $I\mathcal{X}$ is $II\mathcal{X}$. Apply the first CSM formula to $I\mathcal{X}$, push the resulting class from IX to X , and use naturality together with $c_{\text{str}}(X) = c^{SM}(\Phi_X)$. The composite map is $\pi \circ q$, which gives the stated double-inertia formula.

Consequences, difficulty, and scope

Integrating the second formula expresses the stringy Euler characteristic as $\int_{II\mathcal{X}} c_{\text{top}}(T_{II\mathcal{X}})$; the paper also obtains a stringy analogue of the Chern number $c_1 c_{\text{top}-1}$. Under the additional assumptions that X is Gorenstein, π is birational, and $K_{\mathcal{X}} = \pi^* K_X$, Proposition 4.6 translates the result into a formula for orbifold Hodge numbers and proves the cited conjectural identity.

The technically sensitive point is not a formal Gauss–Bonnet argument: it is matching stacky and equivariant pushforwards so that every centralizer weight is correct before applying MacPherson naturality. The main formulas require a smooth separated finite-type Deligne–Mumford stack over $\mathbf{C}$ with projective coarse space; the later Hodge-number statement uses the stronger assumptions just listed.