

S-integral preperiodic points for dynamical systems over number fields

Clayton Petsche

arXiv:0709.3879v2 (2007), 11 pages

<http://arxiv.org/abs/0709.3879v2>

One-sentence summary. For a rational map on the projective line over a number field, Petsche proves Ih’s predicted finiteness of S -integral preperiodic points relative to a non-preperiodic point, under a place-by-place Fatou-set hypothesis on that point.

The Diophantine question and the local hypothesis

Let K be a number field, let S be a finite set of places containing the archimedean ones, and let $\phi : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be defined over K with $\deg \phi \geq 2$. Points $P, Q \in \mathbf{P}^1(\overline{K})$ are S -integral relative to one another when their Zariski closures have disjoint reduction outside S . Ih’s conjecture says that, if P is not preperiodic, only finitely many preperiodic Q are S -integral relative to P .

For a place v , write $F_v(\phi)$ for the Fatou set in $\mathbf{P}^1(\mathbf{C}_v)$. The point P is *totally Fatou at v* if every K -embedding of $K(P)$ into $\mathbf{C}_v$ sends P into $F_v(\phi)$.

Main theorem (Petsche, Theorem 1). *Under the hypotheses above, assume that P is non-preperiodic and totally Fatou at every place of K . Then the set of ϕ -preperiodic points in $\mathbf{P}^1(\overline{K})$ that are S -integral relative to P is finite.*

This applies to arbitrary rational maps, not only maps induced by algebraic groups, but exchanges that generality for a strong local condition on P .

Three ingredients

1. **No accumulation in the Fatou set.** Proposition 2 proves that the preperiodic locus has no limit point in $F_v(\phi)$. Archimedeanly, finiteness of Fatou periodic points and equicontinuity control their backward orbits. Non-archimedeanly, Rivera–Letelier’s classification divides the Fatou set into wandering discs, attracting basins, and the quasiperiodicity domain; each case rules out an accumulation of preperiodic points. This is where the totally Fatou hypothesis later removes the logarithmic poles of the test function from the relevant discrete set.
2. **Canonical local energy.** The dynamical Arakelov–Green functions $g_{\phi,v}$ satisfy

$$\int g_{\phi,v}(R, Q) d\mu_{\phi,v}(R) = 0, \quad \widehat{h}_{\phi}(P) + \widehat{h}_{\phi}(Q) = \frac{1}{[L : K]} \sum_v \sum_{\sigma: L \hookrightarrow \mathbf{C}_v} g_{\phi,v}(\sigma P, \sigma Q),$$

for distinct P, Q contained in a finite extension L/K . At a good-reduction non-archimedean place, $g_{\phi,v}(P, Q) = -\log \delta_v(P, Q)$; it is therefore zero when the two points have distinct reduction. Moreover $\text{supp}(\mu_{\phi,v}) \cap F_v(\phi) = \emptyset$.

3. **Equidistribution of small points.** A sequence of finite Galois-stable sets whose cardinalities tend to infinity and whose average canonical heights tend to zero equidistributes toward $\mu_{\phi,v}$ on the Berkovich projective line. Preperiodic points have canonical height zero, so any hypothetical infinite family supplies exactly such a sequence.

Proof roadmap: one global quantity, two limits

Assume infinitely many admissible preperiodic points exist, and group them into Galois-stable sets Z_n with $|Z_n| \rightarrow \infty$. Let $\mathcal{P}$ be the set of conjugates of P , fix embeddings $\epsilon_v : \bar{K} \hookrightarrow \mathbf{C}_v$, and put

$$\Gamma_v(n) = \frac{1}{|Z_n||\mathcal{P}|} \sum_{Z \in Z_n} \sum_{Q \in \mathcal{P}} g_{\phi,v}(\epsilon_v Z, \epsilon_v Q), \quad \Gamma(n) = \sum_v \Gamma_v(n).$$

The global height identity and $\widehat{h}_\phi(Z) = 0$ give the exact equality

$$\Gamma(n) = \widehat{h}_\phi(P) > 0 \quad \text{for every } n.$$

Enlarge S to contain every bad-reduction place. For $v \notin S$, S -integrality gives distinct reduction, hence $\Gamma_v(n) = 0$. For $v \in S$, the desired test function is

$$F_v(R) = \frac{1}{|\mathcal{P}|} \sum_{Q \in \mathcal{P}} g_{\phi,v}(R, \epsilon_v Q).$$

It has logarithmic poles at the conjugates of P , so equidistribution does not apply directly. Those poles lie in the Fatou set, however: Proposition 2 separates them from all preperiodic points, while the support of $\mu_{\phi,v}$ also avoids them. Thus F_v can be replaced by a bounded continuous F_v^* agreeing with it both on all preperiodic points and on $\text{supp}(\mu_{\phi,v})$. Orthogonality and equidistribution then yield $\Gamma_v(n) \rightarrow 0$. Since S is finite, $\Gamma(n) \rightarrow 0$, contradicting the fixed positive value above.

Example, scope, and the hard point

For $K = \mathbf{Q}$, $S = \{\infty\}$, $\phi(z) = z^2/(z^2 + 4z + 1)$, and $P = \infty$, the hypotheses hold and the theorem says that only finitely many algebraic integers are preperiodic. The proof's delicate step is the simultaneous truncation of every singular local test function without changing either its values on the preperiodic sets or its integral against the canonical measure.

The totally Fatou condition is essential to this *local* method, not claimed to be necessary for Ih's conjecture. When a conjugate of P lies in a nonempty Julia set, it can be approached by preperiodic points, so the separation used in the truncation fails and genuinely global Diophantine input should be expected.