

Rationality of the moduli spaces of plane curves of sufficiently large degree

Christian Böhning, Hans-Christian Graf v. Bothmer

arXiv:0804.1503v1 (2008), 18 pages; DOI: 10.1007/s00222-009-0214-6

<http://arxiv.org/abs/0804.1503v1>

One-sentence summary. Böhning and von Bothmer prove that the moduli space of complex plane curves of degree d is rational for all sufficiently large d , with explicit bounds in every residue class modulo three.

The quotient and the theorem

Put $V(d) = \text{Sym}^d(\mathbf{C}^3)^\vee$. A degree- d plane curve is represented by a point of $\mathbf{P}V(d)$, and projective change of coordinates gives the coarse quotient

$$\text{Hyp}(d, 2) = \mathbf{P}V(d) / \text{PGL}_3(\mathbf{C}).$$

Rationality means that this quotient is birational to a projective space; it is stronger than stable rationality.

Main theorem (Böhning–von Bothmer, Theorem 1.1). *The moduli space $\text{Hyp}(d, 2)$ is rational for all sufficiently large d . More precisely, it is rational when*

$d \bmod 3$	bound
0	$d \geq 1821$,
1	$d \geq 37$,
2	$d \geq 65$.

The divisible-by-three case is an earlier theorem of Katsylo, imported without improving its bound. The paper proves the other two infinite series by reducing high-degree plane curves to fixed low-degree quotients.

Low-degree covariants

Let $G = \text{SL}_3(\mathbf{C})$. For $d = 3n + 1$ the symbolic method constructs a G -equivariant covariant

$$S_d : V(d) \longrightarrow V(4),$$

and for $d = 3n + 2$ a covariant $T_d : V(d) \rightarrow V(8)$. Both are homogeneous of degree four in the coefficients of the input form. Symbolically, if

$$I(\alpha, \beta, \gamma, \delta) = (\alpha\beta\gamma)(\alpha\beta\delta)(\alpha\gamma\delta)(\beta\gamma\delta),$$

then $S_d = I^n \alpha_x \beta_x \gamma_x \delta_x$, while T_d replaces the four final linear factors by their squares. Thus a high-degree quotient is fibered equivariantly over quartics or octics.

The key geometric choice is

$$L_S = x_1^{2n+3} \mathbf{C}[x_1, x_2, x_3]_{n-2} \quad (d = 3n + 1),$$

and analogously $L_T = x_1^{2n+5} \mathbf{C}[x_1, x_2, x_3]_{n-3}$ for $d = 3n + 2$. Proposition 4.1 uses the two torus weights of G to show that $\mathbf{P}L_S$ (respectively $\mathbf{P}L_T$) lies in the covariant's base scheme with full triple structure. Because the covariant has coefficient degree four, its restriction to every $\mathbf{P}(L_S + \mathbf{C}g)$ is linear. This converts nonlinear invariant theory into linear algebra on a moving fiber.

The hard point: fiberwise surjectivity

Proposition 5.3 supplies a g for which

$$S_d|_{\mathbf{P}(L_S + \mathbf{C}g)} : \mathbf{P}(L_S + \mathbf{C}g) \longrightarrow \mathbf{P}V(4)$$

is surjective when $d = 3n + 1 \geq 37$; the corresponding T_d restriction is surjective when $d = 3n + 2 \geq 65$.

The proof first uses Lagrange interpolation to express a one-parameter family $f(c) \in L_S$ or L_T as a sum of d th powers of linear forms. Writing $S_d(f(c) + g) = \sum Q_i c^i$ in the quartic branch, or the analogous T_d expansion in the octic branch, linearity puts every $[Q_i]$ in the image, and the high coefficients become independent of the interpolation points. Their dependence on n has the controlled form $\sum_\nu \rho_\nu^n P_\nu(n)$.

For $d = 3n + 1$, 22 constructed quartics yield a 15×22 coefficient matrix. Modulo 11 it is periodic in n with period 110, and the cited computer-algebra calculation verifies rank 15 throughout; the inequality $n \geq 12$ gives $d \geq 37$. For $d = 3n + 2$, 57 octics give a 45×57 matrix, periodic modulo 19 with period 342 and of rank 45; $n \geq 21$ gives $d \geq 65$. This finite-field rank verification is the proof's most computationally sensitive input.

From fibers to rationality

Let π_{L_S} denote rational projection away from $\mathbf{P}L_S$ onto $\mathbf{P}(V(d)/L_S)$, and write $\bar{g} = \pi_{L_S}(g)$. Consider the incidence variety

$$X = \{(g, \bar{g}, f) : \pi_{L_S}(g) = \bar{g}, S_d(g) = f\}.$$

Fiberwise surjectivity makes the projection from X to $\mathbf{P}(V(d)/L_S) \times \mathbf{P}V(4)$ dominant. Since the restrictions along L_S are linear, X is birational to a vector bundle over that rational base.

On the very stable locus, the quotient $\mathbf{P}V(4) \rightarrow \mathbf{P}V(4)/\mathrm{PGL}_3$ is a principal PGL_3 -bundle in the Zariski topology and hence has a generic section. Pulling the vector-bundle picture back along this section gives

$$\mathbf{P}V(d)/\mathrm{PGL}_3 \sim_{\mathrm{bir}} (\mathbf{P}V(4)/\mathrm{PGL}_3) \times \mathbf{P}^N, \quad N = \dim V(d) - \dim V(4).$$

The quartic quotient becomes rational after multiplying by $\mathbf{P}^8$, and $N \geq 8$ in the stated range. The octic case is identical, using the cited level-eight stable rationality of the octic quotient.

Toy model and caveats

The triple-base-locus phenomenon is the useful toy model: a degree-four map restricted to $L + \mathbf{C}g$ loses its first three nonlinear layers and becomes linear. Surjectivity then makes a general fiber look like an affine space varying over a rational base. The full proof must additionally make this construction equivariant and descend it through the quotient.

The theorem concerns birational coarse moduli over $\mathbf{C}$, not an explicit normal form for every curve. The bounds are not claimed optimal; the paper specifically leaves small cases such as plane sextics open. Its rank checks rely on the authors' referenced Macaulay2 scripts, while the $d \equiv 0 \pmod{3}$ range relies on Katsylo's separate result.