

Rational Periodic Points for Degree Two Polynomial Morphisms on Projective Space

Benjamin Hutz

arXiv:0811.3225v2 (2009), 12 pages

<http://arxiv.org/abs/0811.3225v2>

One-sentence summary. Hutz constructs infinite families of quadratic polynomial morphisms of projective space with long rational periodic orbits, then combines relatively prime periods to obtain super-polynomial growth with the dimension.

Polynomial dynamics and the theorem

A degree-two *polynomial map* of $\mathbf{P}^N$ has the form

$$\phi = [\phi_0(x_0, \dots, x_N), \dots, \phi_{N-1}(x_0, \dots, x_N), x_N^2],$$

where the first N entries are quadratic forms over $\mathbf{Q}$. It is a morphism precisely when these homogeneous coordinates have no common nonzero zero. A point has primitive period n if $\phi^n(P) = P$ and no smaller positive iterate returns it.

Put

$$T_2 = 7, \quad T_N = \binom{N+2}{2} + \left\lfloor \frac{N-1}{2} \right\rfloor \quad (N \geq 3).$$

Theorems 1 and 2 give an infinite family, of dimension at least N , of degree-two polynomial morphisms $\phi : \mathbf{P}^N \rightarrow \mathbf{P}^N$ having a $\mathbf{Q}$ -rational point of primitive period T_N . Closing the orbit earlier also realizes every period from 1 through $\binom{N+2}{2}$. The asymptotic conclusion is stronger.

Main theorem (Hutz, Theorem 3). *For every fixed positive integer k , all sufficiently large N admit a degree-two polynomial morphism of $\mathbf{P}^N$ over $\mathbf{Q}$ with a $\mathbf{Q}$ -rational periodic point of primitive period greater than $c(k)N^k$, for some constant $c(k) > 0$ depending only on k .*

Thus, unlike the conjectural uniform behavior for quadratic polynomials on $\mathbf{P}^1$, attainable rational periods in higher dimension cannot be bounded by any fixed polynomial in N .

Ingredient I: a triangular orbit prescription

Write

$$\phi_i = \sum_{0 \leq j \leq \ell \leq N} c_i(j, \ell) x_j x_\ell, \quad \phi_N = x_N^2.$$

Starting at $P = [0 : \dots : 0 : 1]$, Lemma 1 prescribes a chain of 0–1 points by turning on consecutive blocks of coordinates. The order is chosen so that each next image is linear in one still-unused coefficient $c_i(j, \ell)$. This spends the off-diagonal coefficients first and leaves the diagonal blocks $c_i(j, j)$ available after roughly $\binom{N+2}{2}$ steps.

Each remaining diagonal block can usually add two orbit points. Along a coordinate slice, every ϕ_i is a quadratic polynomial in one variable; three already prescribed input-output pairs therefore determine the next image. A fresh diagonal coefficient then prescribes the image after that. Excluding finitely many parameter values keeps the new points distinct, so the return has primitive, not merely divisible, period. This interpolation mechanism supplies the extra $\lfloor (N-1)/2 \rfloor$ steps in T_N .

Ingredient II: maps versus morphisms

The orbit construction initially gives rational maps. To exclude a base point, the paper uses the Macaulay resultant of the $N+1$ coordinate quadrics. Its vanishing is exactly the condition that they have a common nontrivial zero. The proof partitions degree- $(N+2)$ monomials into sets S_i according to the last square dividing them and selects a maximal minor of the Macaulay matrix. The coefficient pattern forced by the orbit construction makes this minor a nonzero polynomial in the free parameters. Consequently the resultant does not vanish identically on the family, and infinitely many rational parameter choices are genuine morphisms. This is the technically delicate gate: a long formal orbit is useful only after indeterminacy has been removed.

Ingredient III: multiply coprime periods

Lemma 2 is the amplification step. If polynomial morphisms on $\mathbf{P}^a$ and $\mathbf{P}^b$ have points of primitive periods m and n , their product on affine charts, followed by homogenization, gives a polynomial morphism on $\mathbf{P}^{a+b}$ with primitive period $\text{lcm}(m, n)$. The two coordinate blocks share only the homogenizing variable, so no new common zero is introduced.

Fix s and take blocks of dimension $M \sim N/s$. The first construction realizes every prime

$$p \leq \binom{M+2}{2} \sim \frac{N^2}{2s^2}.$$

Choose s distinct primes in a short interval below this bound. Their least common multiple is their product, and for any fixed $\varepsilon > 0$ the proof obtains

$$p_1 \cdots p_s \geq \frac{(1-\varepsilon)^s}{2^s s^{2s}} N^{2s}.$$

Taking $2s > k$ proves Theorem 3; a period-one block fills any leftover dimension.

Toy model, difficulty, and scope

On $\mathbf{P}^1$, the quadratic $f(x) = ax^2 + bx + c$ is determined by three prescribed values. Choosing $0 \mapsto 1 \mapsto 2 \mapsto 0$ yields $f(x) = -\frac{3}{2}x^2 + \frac{5}{2}x + 1$, a rational 3-cycle. The higher-dimensional construction is a triangular version of this interpolation, with many coefficient blocks and enough coordinate slices to reuse the quadratic three-value rule.

The results are existential lower bounds, not upper bounds or a classification of possible periods. The paper's separate examples in dimensions 2, 3, 4 have periods 9, 24, 72, beyond the uniform family, but the omitted resultant calculations for those examples are verified only by explicit computation. The construction is specific to quadratic polynomial morphisms over $\mathbf{Q}$.