

Deformations of canonical pairs and Fano varieties

Tommaso de Fernex, Christopher D. Hacon

arXiv:0901.0389v3 (2009), 26 pages

<http://arxiv.org/abs/0901.0389v3>

One-sentence summary. De Fernex and Hacon prove that every simplicial complex toric Fano variety with at most terminal singularities is rigid under small projective flat deformations, by transporting its class group and polynomial Cox ring through the family.

Rigidity and the exact hypotheses

Work over $\mathbf{C}$. A normal projective variety X_0 is Fano when $-K_{X_0}$ is $\mathbf{Q}$ -Cartier and ample. For a toric variety, simpliciality of the fan is equivalent to $\mathbf{Q}$ -factoriality. The terminal condition is the strict discrepancy condition used in the minimal model program.

Main theorem (de Fernex–Hacon, Theorems 1.4 and 7.1). *Every simplicial toric Fano variety with at most terminal singularities is rigid under small projective flat deformations. Hence, if it is singular, it is not smoothable by such a deformation.*

Concretely, for a projective flat family $f : \mathcal{X} \rightarrow T$ over a smooth pointed curve with central fiber X_0 , after shrinking near the marked point (and making the finite base changes used in the proof), the nearby fibers are isomorphic to X_0 and the deformation is locally trivial. This is striking because singular toric varieties often occur as special fibers of useful degenerations; the theorem identifies a sharp rigid regime.

This digest focuses on Theorem 7.1. Other principal results, not developed here, include the adjoint extension theorem in its full generality, the constancy of nef values and movable cones, and preservation of the complete Mori chamber decomposition for Fano varieties in dimension at most three, for 1-canonical fourfolds, and in the toric case (Theorems 5.7, 6.8, and 6.9 and their introductory forms).

First ingredient: classes survive the family

Corollary 3.2 and Propositions 3.5 and 3.8 show that, after shrinking T , the total space and all fibers remain terminal, $\mathbf{Q}$ -factorial, and Fano. Proposition 6.5 then makes the Néron–Severi spaces constant after killing finite monodromy. Numerical divisor classes alone are insufficient for a Cox ring, so Lemma 7.2 proves the integral refinement

$$\mathrm{Cl}(\mathcal{X}/T) \xrightarrow{\sim} \mathrm{Cl}(X_0)$$

after a suitable base change.

The hard part of that lemma cuts the family by general hyperplanes to a smooth family of surfaces $S \rightarrow T$. Terminal singularities have codimension at least three, while $H^1(S_t, \mathcal{O}_{S_t}) = 0$ makes $\mathrm{Pic}(S_t) \hookrightarrow H^2(S_t, \mathbf{Z})$ have torsion-free cokernel. Finite monodromy is removed, finitely many generators of $\mathrm{Cl}(X_0)$ are extended using algebraic families of divisors, and Grothendieck–Lefschetz restriction transfers the resulting isomorphism back from S to $\mathcal{X}$.

Second ingredient: every graded section extends

Theorem 4.5 is the paper’s extension theorem for adjoint Weil divisors. In the Fano situation, the relative ampleness of $-K_{\mathcal{X}}$ lets it be applied to each class $[A] \in \mathrm{Cl}(\mathcal{X}/T)$, giving

$$H^0(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(A)) \rightarrow H^0(X_t, \mathcal{O}_{X_t}(A)) \quad \text{for every } t.$$

Its proof runs a relative minimal model program that restricts step by step to the central fiber, then uses a short exact restriction sequence and vanishing on the resulting minimal model. This is where the Fano positivity and terminal/ $\mathbf{Q}$ -factorial hypotheses enter the rigidity argument rather than merely simplifying notation.

Third ingredient: deform the Cox polynomial ring

The total coordinate ring of the toric central fiber is

$$R_0 = \bigoplus_{[D] \in \text{Cl}(X_0)} H^0(X_0, \mathcal{O}_{X_0}(D)) = \mathbf{C}[x_{0,1}, \dots, x_{0,r}],$$

with one variable for each ray of the fan and its natural class-group grading. Via Lemma 7.2 define the analogous graded ring

$$R_t = \bigoplus_{[A] \in \text{Cl}(\mathcal{X}/T)} H^0(X_t, \mathcal{O}_{X_t}(A)).$$

The extension theorem lifts each $x_{0,i}$ to $x_{t,i}$ and produces

$$\Phi_t : \mathbf{C}[x_1, \dots, x_r] \longrightarrow R_t, \quad x_i \longmapsto x_{t,i}.$$

To prove that Φ_t remains an isomorphism, choose an ample divisor to turn the class-group grading into a positive integer grading and let m and M be the smallest and largest degrees of the central variables. The sums of section spaces of degree at most d form trivial vector bundles on T . For each ordinary polynomial degree e , injectivity of the bounded map into classes of degree at most Me and surjectivity onto classes of degree at most me are open conditions. Both hold at $t = 0$, hence they hold simultaneously for every bound at very general t . Thus $R_t \cong R_0$ for infinitely many nearby fibers.

Finally, for a relatively ample divisor L the graded isomorphism respects the section subrings, so

$$X_t = \text{Proj} \bigoplus_{q \geq 0} H^0(X_t, qL) \cong \text{Proj} \bigoplus_{q \geq 0} H^0(X_0, qL) = X_0.$$

This recovers the variety, not merely its numerical cones, and completes the local-rigidity argument.

Toy model, hardest point, and limits

For $X_0 = \mathbf{P}^n$, the Cox ring is the standard polynomial ring $\mathbf{C}[x_0, \dots, x_n]$ and $X_0 = \text{Proj } R_0$; preserving the graded ring visibly preserves projective space. The full theorem must first preserve a possibly torsion-bearing class group and every graded piece across a singular family. That class-group/extension bridge is the proof's hardest point.

All hypotheses matter. The paper gives failures for weak or log Fano toric varieties, for canonical singularities that are not terminal, and for terminal toric Fano varieties that are not simplicial. The conclusion concerns small projective flat deformations over $\mathbf{C}$, not arbitrary analytic or nonprojective deformations.