

Moduli spaces for families of rational maps on the projective line

Michelle Manes

arXiv:0902.1813v2 (2009), 47 pages

<http://arxiv.org/abs/0902.1813v2>

One-sentence summary. Manes proves that quadratic rational maps with a marked formal-period point form a geometrically irreducible moduli space for every level $N > 1$, while nontrivial automorphisms create special loci with systematic reducibility.

Dynatomic level structure

Let $\phi = [F_1 : G_1] : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ have degree d , and write $\phi^k = [F_k : G_k]$. Its homogeneous dynatomic polynomial is

$$\Phi_{N,\phi}^*(x, y) = \prod_{k|N} (yF_k(x, y) - xG_k(x, y))^{\mu(N/k)}.$$

This Möbius product is a polynomial. Its roots have *formal period* N : every point of primitive period N occurs, but exceptional points of smaller primitive period can occur with multiplicity. With Rat_d the space of degree- d maps, set

$$\text{Rat}_d(N) = \{(P, \phi) : \Phi_{N,\phi}^*(P) = 0\}, \quad M_d(N) = \text{Rat}_d(N) / \text{PGL}_2.$$

Proposition 3.2 constructs these as schemes over $\mathbf{Z}$; the quotient records a conjugacy class together with its marked formal-period point.

Main theorem (Manes, Theorem 4.5). *For every integer $N > 1$, the level moduli space $M_2(N)$ is geometrically irreducible.*

Thus irreducibility is not peculiar to the polynomial line $z \mapsto z^2 + c$: it holds for the full two-dimensional moduli space of quadratic rational maps. Consequently, Bertini gives generic irreducibility for period- N curves cut out by one-parameter families (Corollary 4.6).

Ingredient I: a two-parameter normal form

Away from the exceptional class $(3, 3) \in M_2$, the family

$$\phi_{a,b}[x : y] = [x^2 + axy : bxy + y^2], \quad ab \neq 1,$$

maps surjectively and generically six-to-one to $M_2 \setminus \{(3, 3)\}$. The parameters a, b are multipliers of two fixed points. Hence the universal equation $\Phi_{N,\phi_{a,b}}^* = 0$ is a hypersurface over this normal-form chart and maps to $M_2(N)$.

Lemma 4.2 computes the extreme coefficients:

$$[x^{\nu_2(N)}] \Phi_{N,\phi_{a,b}}^* = C_N(b), \quad [y^{\nu_2(N)}] \Phi_{N,\phi_{a,b}}^* = C_N(a),$$

where C_N is the N th cyclotomic polynomial. This calculation is the control needed for a specialization argument: it prevents a putative factor from disappearing when a parameter is set to zero.

Ingredient II: specialize without losing a factor

Proposition 4.3 proves the universal dynatomic polynomial irreducible for $N > 1$. If it factored nontrivially as AB with both factors of positive x -degree, their leading coefficients would multiply to $C_N(b)$. Because $C_N(0) = 1$ for $N > 1$, specializing $b = 0$ preserves the positive x -degree of both factors. But

$$\phi_{a,0}(z) = z^2 + az,$$

and Morton's irreducibility theorem (Lemma 4.4) says that this specialized dynatomic polynomial is irreducible. The contradiction proves that the normal-form hypersurface is irreducible. The hardest point is precisely the cyclotomic leading-coefficient check: irreducibility of a special fiber would say nothing if an alleged factor collapsed under specialization.

Ingredient III: descend to moduli

For $N > 1$ the normal-form hypersurface surjects onto $M_2(N)$ after removing only the finite fiber above $(3, 3)$; horizontally the map is generically six-to-one and vertically it has the expected dynatomic degree $\nu_2(N)$. The image of the geometrically irreducible hypersurface is therefore irreducible, and taking the finite closure gives Theorem 4.5. This last step is why the proof works on conjugacy classes rather than only on a chosen coordinate normal form.

A toy boundary case

At level one the same family has

$$\Phi_{1, \phi_{a,b}}^*(x, y) = xy(x(1-b) + y(1-a)).$$

The factors x and y are the two fixed points deliberately placed at 0 and ∞ when the normal form is chosen; the third factor supplies the remaining fixed point. This explains both why the proof must assume $N > 1$ and why formal-period level structure is the right invariant object.

The automorphism contrast and scope

The paper's other principal theme is deliberately nongeneric. If $h \in \text{Aut}(\phi)$ has prime order p , an h -tuned dynatomic factor detects points for which $h(P)$ lies on the same ϕ -orbit as P . Sections 6–7 prove that this factor is a genuine, proper divisor of Φ_{pN}^* for all but finitely many relevant N . Accordingly, Corollary 7.10 shows that $M_d(pN, C_p)$ is geometrically reducible for all but finitely many N in characteristic zero, and for all but finitely many prime N over an arbitrary field. Its components separate the cases where P and $h(P)$ lie on one orbit or on distinct orbits.

The digest proves neither that every formal-period point has primitive period N nor that every one-parameter family gives an irreducible curve. It also omits the tuned multiplicity calculations, the pure-power factorizations, and the final analysis of the order-two symmetry curve; those results refine the exceptional automorphism locus rather than the global irreducibility theorem explained here.