

Globally F -regular and log Fano varieties

Karl E. Schwede, Karen E. Smith

arXiv:0905.0404v2 (2009), 31 pages

<http://arxiv.org/abs/0905.0404v2>

One-sentence summary. Schwede and Smith convert Frobenius splittings into effective boundaries, showing that global F -regularity is the prime-characteristic shadow of log Fano geometry and proving the converse after reduction from characteristic zero.

The dictionary and the exact statements

Let X be normal and F -finite of characteristic $p > 0$, and let $B \geq 0$ be a $\mathbf{Q}$ -divisor. The pair (X, B) is globally F -regular if, for every effective divisor D , some Frobenius power makes the natural map

$$\mathcal{O}_X \longrightarrow F_*^e \mathcal{O}_X(\lceil (p^e - 1)B \rceil + D)$$

split as an $\mathcal{O}_X$ -module map. A pair (X, Δ) is log Fano when it is Kawamata log terminal (klt) and $-(K_X + \Delta)$ is ample.

Main theorem (Schwede–Smith, Theorems 4.3, 1.1, and 5.1). *Suppose X is quasiprojective over an F -finite local ring with a dualizing complex. If (X, B) is globally F -regular, there is an effective $\mathbf{Q}$ -divisor Δ such that $(X, B + \Delta)$ remains globally F -regular and $K_X + B + \Delta$ is anti-ample; the coefficient denominators may be chosen prime to p . Consequently every globally F -regular variety in prime characteristic is log Fano.*

Conversely, if X is normal projective over a characteristic-zero field and (X, Δ) is klt with $K_X + \Delta$ anti-ample, then the pair has globally F -regular type.

Here “type” means that suitable models have globally F -regular fibers for almost all reductions modulo p ; it does not assert global F -regularity in characteristic zero itself.

Ingredient I: dualize a Frobenius splitting

Given a splitting

$$\mathcal{O}_X \longrightarrow F_*^e \mathcal{O}_X(D) \longrightarrow \mathcal{O}_X,$$

Frobenius duality and a canonical twist produce another splitting

$$\mathcal{O}_X \longleftarrow F_*^e \mathcal{O}_X((1 - p^e)K_X - D) \longleftarrow \mathcal{O}_X.$$

The image of 1 in the middle sheaf cuts out an effective divisor $D' \sim (1 - p^e)K_X - D$. This turns an abstract retraction into a concrete boundary with a controlled canonical class.

For a globally sharply F -split pair (X, B) , take $D = (p^e - 1)B$. Then $\Delta_1 = D'/(p^e - 1)$ keeps the pair sharply split and

$$K_X + B + \Delta_1 \sim_{\mathbf{Q}} 0.$$

For a globally F -regular pair, insert a very ample effective divisor H into the splitting. The same dual construction gives $\Delta_2 \geq 0$ with

$$K_X + B + \Delta_2 \sim_{\mathbf{Q}} -\frac{1}{p^f - 1}H,$$

which is anti-ample.

Ingredient II: retain regularity while gaining positivity

The second boundary is initially only sharply F -split. Lemma 4.9 is the technical bridge: suitable rational convex combinations of sharply split boundaries stay sharply split, while combinations starting from a globally F -regular boundary can be chosen globally F -regular. After arranging the support of Δ_1 inside that of H , two applications of the lemma give positive $\epsilon_1 + \epsilon_2 = 1$ such that

$$\Delta = \epsilon_1 \Delta_1 + \epsilon_2 \Delta_2$$

makes $(X, B + \Delta)$ globally F -regular. Its canonical class is the same convex combination of a $\mathbf{Q}$ -trivial and an anti-ample class, hence is anti-ample. Keeping the exponents in the splitting formulas also keeps all boundary denominators prime to p . Controlling the singularity type through this interpolation, rather than merely manufacturing an effective divisor, is the proof’s hardest point.

Ingredient III: pass from F -regular to log Fano

Global F -regularity implies local F -regularity. Because the constructed $K_X + B + \Delta$ is $\mathbf{Q}$ -Cartier, the Hara–Watanabe comparison used in Theorem 4.4 turns local F -regularity into klt singularities. Anti-ampleness then gives the log Fano conclusion. Notice that the boundary is essential: the theorem says more than that $-K_X$ is big, and it does not require X to have been $\mathbf{Q}$ -Gorenstein before Δ is constructed.

The converse: test the vertex of an affine cone

For an ample invertible sheaf $\mathcal{L}$, let

$$S(X, \mathcal{L}) = \bigoplus_{n \geq 0} H^0(X, \mathcal{L}^n).$$

Proposition 5.4 says that a characteristic-zero pair is log Fano exactly when one can choose such a section ring whose affine cone $(\operatorname{Spec} S, \Delta_S)$ is klt. Blowing up the vertex identifies the exceptional divisor with X ; adjunction shows that the discrepancy at the vertex records the sign of $K_X + \Delta$. A klt cone has globally F -regular type by the local characteristic- p comparison. Proposition 5.3 then identifies global F -regularity of the cone with global F -regularity of the projective pair, completing Theorem 5.1.

As a toy case, $\mathbf{P}^n$ already has $-K_{\mathbf{P}^n} = (n+1)H$ ample, so the boundary can be 0 and the cone is affine space. The general proof replaces this transparent polynomial cone by a singular section ring and uses Frobenius duality to create the missing boundary; the toy case does not reflect the non- $\mathbf{Q}$ -Gorenstein difficulty.

Further contributions and limits

The same construction shows that a globally sharply F -split variety admits a log Calabi–Yau boundary: $K_X + \Delta \sim_{\mathbf{Q}} 0$ and the pair is log canonical. Section 6 also proves a Kawamata–Viehweg-type vanishing theorem for globally F -regular pairs. These are principal companion results, but their full proofs are omitted here.

The converse is not a statement in each small characteristic: the paper’s smooth cubic surface example is log Fano but not even F -split in characteristic 2. Nor does the paper prove the conjectural converse for log Calabi–Yau pairs; it records dense global sharp F -split type as an expectation. All claims above concern normal pairs under the stated F -finiteness, projectivity, and duality hypotheses.