

Preperiodic points and unlikely intersections

Matthew Baker, Laura DeMarco

arXiv:0911.0918v3 (2010), 26 pages

<http://arxiv.org/abs/0911.0918v3>

One-sentence summary. Baker and DeMarco prove that two fixed complex points are simultaneously preperiodic for infinitely many maps $z^d + c$ exactly when their d th powers agree, by making parameter-space potential theory interact with adelic equidistribution.

The unlikely-intersection theorem

Fix $d \geq 2$, put $f_c(z) = z^d + c$, and write

$$\text{Preper}(a) = \{c \in \mathbf{C} : a \text{ is preperiodic for } f_c\}.$$

The condition on one marked point is sparse but infinite; the theorem says that two independent such conditions have only finite intersection.

Main theorem (Baker–DeMarco, Theorem 1.1). *For fixed $a, b \in \mathbf{C}$, the set of parameters $c \in \mathbf{C}$ for which both a and b are preperiodic for f_c is infinite if and only if $a^d = b^d$.*

The easy direction is already dynamical: $a^d = b^d$ gives $f_c(a) = f_c(b)$ for every c , and Lemma 3.5 shows that $\text{Preper}(a)$ is infinite. The reverse direction is the arithmetic heart of the paper.

Ingredient I: a Mandelbrot set for each marked point

Let G_c be the escape-rate Green function of f_c and define

$$M_a = \{c : \sup_n |f_c^n(a)| < \infty\}, \quad G_a(c) = G_c(a^d + c).$$

Proposition 3.3 proves that M_a is compact and full, that G_a is its Green function, that its logarithmic capacity is one, and that its equilibrium measure μ_a is supported on ∂M_a . Near infinity the Böttcher coordinate ϕ_c gives the normalized uniformizer

$$\Phi_a(c) = \phi_c(a^d + c), \quad G_a(c) = \log |\Phi_a(c)|, \quad \Phi'_a(\infty) = 1.$$

Montel’s theorem makes preperiodic parameters dense along the relevant boundary. Theorem 3.4 then packages the complex-analytic rigidity:

$$M_a = M_b \iff \mu_a = \mu_b \iff a^d = b^d \iff \text{Preper}(a) = \text{Preper}(b).$$

The nonformal step at the center uses fullness to recover M_a from the support of μ_a , then uses $\Phi_a = \Phi_b$ and injectivity of ϕ_c near infinity to force $a^d + c = b^d + c$.

Ingredient II: algebraic parameters equidistribute

Assume first that $a, b \in \overline{\mathbf{Q}}$ and that distinct c_n make both points preperiodic. An orbit relation $f_{c_n}^\ell(a) = f_{c_n}^m(b)$ shows that each c_n is algebraic. Preperiodicity also gives height zero for the adelic set attached to M_a , and the same remains true for every Galois conjugate of c_n . Corollary 2.10 therefore sends the uniform measures on those conjugates to μ_a . The identical sequence converges to μ_b , so uniqueness of a weak limit yields $\mu_a = \mu_b$; Theorem 3.4 finishes the proof. This is stronger than a mere finiteness argument: Theorem 4.1 says that if $a^d \neq b^d$ are algebraic, then $h_{M_a}(c) + h_{M_b}(c)$ is bounded below by a positive constant outside a finite set.

Ingredient III: the transcendental function-field step

If a is transcendental, the common orbit relations force b and all c_n into the algebraic closure of $k = \overline{\mathbf{Q}}(a) \simeq \overline{\mathbf{Q}}(T)$. For each place v of this product-formula field, the paper constructs a Berkovich generalized Mandelbrot set $M_{a,v}$. Taking growing Galois-stable unions of the conjugates of the c_n , Theorem 2.8 equidistributes their uniform measures toward the equilibrium measure of $M_{a,v}$. Symmetry gives

$$M_{a,v} = M_{b,v} \quad \text{for every place } v.$$

Corollary 3.13, based on Benedetto's function-field theorem, converts membership in all these local filled sets back into preperiodicity. Thus a and b are preperiodic for exactly the same parameters; Theorem 3.4 again gives $a^d = b^d$. Establishing this local-to-global implication in the absence of an archimedean place is the proof's hardest additional point and explains why Berkovich spaces are essential rather than decorative.

A concrete test case

For $d = 2$, $a = 0$, and $b = 1$, the theorem applies because $0^2 \neq 1^2$: only finitely many quadratic parameters can make both marked points preperiodic. The paper checks that $c = 0, -1, -2$ do so. It conjectures that these are the only three, but the equidistribution proof is ineffective and does not bound or enumerate the exceptional parameters. This example captures the unlikely-intersection phenomenon while omitting precisely the arithmetic difficulty responsible for ineffectivity.

Companion result and scope

The same architecture proves Theorem 1.2: two rational functions of degree at least two have infinitely many common preperiodic points exactly when their full preperiodic sets coincide; consequently distinct Julia sets permit only finitely many common preperiodic points. Its separate canonical-height argument is not reproduced here. Nor does the main theorem treat moving marked points or general one-parameter families: the simple conclusion $a^d = b^d$ depends on the uncritical form $z^d + c$.