

Smoothings of Fano varieties with normal crossing singularities

Nikolaos Tziolas

arXiv:1005.0531

<http://arxiv.org/abs/1005.0531v2>

One-sentence summary. Tziolas gives smoothing criteria for normal-crossing Fano varieties and proves that a smoothing with smooth total space exists exactly when the infinitesimal deformation line bundle on the singular locus is trivial.

The deformation line bundle

Work over an algebraically closed field k of characteristic zero. A normal-crossing variety is locally

$$\hat{\mathcal{O}}_{X,P} \simeq k(P)[[x_0, \dots, x_d]]/(x_0 \cdots x_r),$$

and it is Fano here when X is reduced and projective and ω_X^{-1} is ample. Let D be the singular locus and set

$$T_X^1 := \mathcal{E}xt_X^1(\Omega_X, \mathcal{O}_X).$$

For normal crossings, T_X^1 is a line bundle on D measuring first-order changes of the singularities. The variety is *d-semistable* when $T_X^1 \simeq \mathcal{O}_D$. A smoothing is a flat family over a discrete valuation ring with special fibre X and smooth generic fibre; it is *total* when the total space itself is smooth.

Main theorem (Tziolas, Theorems 1.1 and 5.3). *Let X be a normal-crossing Fano variety as above. It is smoothable if any one of the following holds:*

- (1) T_X^1 is globally generated and $H^1(T_X^1) = 0$;
- (2) X has only double-point normal crossings and T_X^1 is globally generated;
- (3) X is *d-semistable*.

Moreover, X admits a total smoothing if and only if it is d-semistable.

Ingredient I: remove the global obstructions

Ordinary deformations have obstruction spaces $H^2(T_X)$ and $H^1(T_X^1)$. Theorem 4.7 proves for every such Fano variety that

$$H^2(T_X) = H^2(X, \mathcal{O}_X) = 0;$$

with a semistable log structure it also proves $H^2(\mathcal{H}om(\Omega_X(\log), \mathcal{O}_X)) = 0$. The hard input is the normalization-strata resolution of Ω_X/τ_X in Theorem 4.4, where $\tau_X \subset \Omega_X$ is the torsion subsheaf. After Serre duality, its terms become differential forms on smooth normalizations of successive singular strata, twisted by ω_X and two-torsion line bundles. The Fano condition makes the dual twists ample, so Akizuki–Kodaira–Nakano vanishing kills the needed cohomology groups inductively.

For double crossings, Theorem 4.10 also kills $H^1(T_X^1)$. Indeed D is a smooth Fano variety, and global generation makes $T_X^1 \otimes \omega_D^{-1}$ ample; Kawamata–Viehweg vanishing gives $H^1(T_X^1) = 0$. Proposition 5.1 then turns these vanishings and global generation into an ordinary smoothing, proving the first two criteria.

Ingredient II: log smoothness globalizes $x_0 \cdots x_r = t$

Kato's criterion (Theorem 3.9) says that X carries a semistable log structure exactly when $T_X^1 \simeq \mathcal{O}_D$. Choose a nowhere-vanishing section of this trivial bundle. Etale locally it is the first-order deformation

$$\frac{k[x_0, \dots, x_d]}{(x_0 \cdots x_r)} \rightsquigarrow \frac{k[t]/(t^2)[x_0, \dots, x_d]}{(x_0 \cdots x_r - t)}.$$

The logarithmic obstruction group vanishes by Theorem 4.7, so Corollary 4.9 lifts this direction through all infinitesimal orders to a formal log deformation over $k[[t]]$. The vanishing of $H^2(\mathcal{O}_X)$ makes that formal deformation effective. Finally the relative cotangent sequence and the nonzero smoothing direction make the map $\mathcal{O}_X \rightarrow T_{\mathcal{X}/\Delta}^1$ surjective; the resulting $\mathcal{E}xt_{\mathcal{X}}^1(\Omega_{\mathcal{X}}, \mathcal{O}_{\mathcal{X}})$ vanishes, and the complete-intersection total space $\mathcal{X}$ is smooth.

This log-deformation step is stronger than ordinary unobstructedness: it preserves the local semistable equation while making the smoothing direction coherent across all branches.

Why triviality is also necessary

Conversely, suppose X is the special fibre of a smoothing with smooth total space $\mathcal{X}$. Dualizing the relative cotangent sequence and restricting to X gives a surjection $\mathcal{O}_D \twoheadrightarrow T_X^1$. Since T_X^1 is a line bundle on D , the smoothing parameter trivializes it. Thus total smoothability forces d -semistability (Proposition 5.1), closing the equivalence.

The double-crossing toy model

For $X = X_1 \cup_D X_2$ with smooth components meeting transversely,

$$T_X^1 \simeq N_{D/X_1} \otimes N_{D/X_2}.$$

Hence d -semistability says that the two normal directions cancel, exactly what is needed to replace the local node $xy = 0$ by $xy = t$. Section 6 glues components along $D \simeq \mathbf{P}^1 \times \mathbf{P}^1$: normal bundles $\mathcal{O}_D(-1, -1)$ and $\mathcal{O}_D(1, 1)$ give $T_X^1 \simeq \mathcal{O}_D$ and a smoothing, whereas gluing two negative sides gives $\mathcal{O}_D(-2, -2)$, no global smoothing direction, and only locally trivial deformations.

Scope and open edge

The Fano hypothesis is decisive: d -semistability is not sufficient for arbitrary normal-crossing varieties. Characteristic zero is also essential to the proof's vanishing theorems. The paper leaves open whether every smoothable normal-crossing Fano has globally generated T_X^1 . It separately asks whether the ordinary deformation functor is smooth for every such Fano; a positive answer would imply the proposed characterization by global generation. The theorem does not address more general singularities or classify the smooth Fano fibres.