

Existence of non-preperiodic algebraic points for a rational self-map of infinite order

Ekaterina Amerik

arXiv:1007.1635

<http://arxiv.org/abs/1007.1635v1>

One-sentence summary. Amerik proves that a dominant rational self-map of infinite order over a number field has an algebraic point with infinite forward orbit, by trapping the dynamics in a good p -adic neighbourhood and analytically interpolating its iterates.

The question beyond canonical heights

Let X be a variety over a number field K and $f: X \dashrightarrow X$ a dominant rational self-map. A point is preperiodic only when every required iterate is defined and its forward orbit is finite. If f is a polarized morphism, canonical heights immediately produce non-preperiodic algebraic points. For a rational map, however, indeterminacy and the lack of a satisfactory canonical height remove that argument.

Main theorem (Amerik, Corollary 9). *If f is not of finite order as a rational self-map, then there exists $x \in X(\overline{\mathbf{Q}})$ that is not preperiodic under f .*

The result is deliberately weaker than the conjectural existence of a Zariski-dense algebraic orbit, but it applies without regularity or polarization hypotheses.

Ingredient I: manufacture an invariant p -adic polydisc

Choose a prime of good reduction for an affine model of (X, f) . Applied to the graph of the reduced map, Hrushovski's finite-field theorem supplies a periodic smooth point $\bar{x}$ whose entire orbit avoids both indeterminacy and ramification (Corollary 2). After a finite extension of K , Hensel lifting gives a point y above $\bar{x}$ and a residue neighbourhood

$$\mathcal{O}_{\mathfrak{p},y} = \{t : x_i(t) \equiv x_i(y) \pmod{\mathfrak{p}}\}.$$

If k is the period of $\bar{x}$, Proposition 3 identifies this neighbourhood with $\mathfrak{p}^n$, makes f^k a bijection of it, keeps every iterate away from the bad loci, and makes $X(\overline{\mathbf{Q}})$ dense in it. Thus the rational dynamics has become everywhere-defined analytic dynamics on one compact p -adic chart.

Ingredient II: interpolate the discrete orbit

After rescaling by a uniformizer r , write the restriction of f^k as a convergent map $F: \mathcal{O}_{\mathfrak{p}}^n \rightarrow \mathcal{O}_{\mathfrak{p}}^n$. Its higher coefficients satisfy the divisibility needed in the Bell–Ghioca–Tucker construction. Since F modulo $\mathfrak{p}$ is an affine automorphism of a finite affine space, Proposition 4 gives an $l > 0$ with

$$F^l(z) \equiv z \pmod{\mathfrak{p}} \quad \text{for every } z.$$

Theorem 7 then interpolates the iterates of a suitable further power $\psi = f^N$: for each starting point ω there is a vector-valued function g , continuous on $\mathbf{Z}_p$ and analytic on a smaller ball $p^a \mathbf{Z}_p$, such that

$$g(0) = \omega, \quad g(p^a i) = \psi^i(\omega) \quad (i \geq 0)$$

after restricting to the relevant congruence class.

The interpolation is built from Mahler series $g_j(z) = \omega_j + \sum_{m \geq 1} b_{jm} \binom{z}{m}$. The paper's Lemma 6 is the technical repair needed over the possibly ramified valuation ring $\mathcal{O}_{\mathfrak{p}}$: the coefficient estimates may not give analyticity on all of $\mathbf{Z}_p$, but they do give it on some $p^a \mathbf{Z}_p$. This passage from integral coefficients to a ramified extension is the proof's most delicate calculation.

Ingredient III: finite orbit forces uniformly bounded period

Suppose $\omega \in \mathcal{O}_{\mathfrak{p},y}$ is preperiodic. Its interpolated orbit is finite, so one vector value of g occurs at infinitely many integers in the compact ball $p^a \mathbf{Z}_p$. Those integers have an accumulation point; the p -adic identity principle forces every coordinate of g to be constant. Consequently $\psi(\omega) = \omega$. Corollary 8 therefore gives one integer N , depending only on the neighbourhood, such that every preperiodic point in it is fixed by f^N .

Now f has infinite order, hence f^N is not the identity rational map. Its fixed locus inside the good neighbourhood is therefore a proper analytic subvariety. It cannot contain the dense set of algebraic points from Proposition 3. Any algebraic point outside it has a well-defined infinite orbit, proving the theorem.

A one-variable model of the rigidity step

The last implication is already visible without the geometry. If an analytic map $g: p^a \mathbf{Z}_p \rightarrow \mathcal{O}_{\mathfrak{p}}$ records an orbit by $g(p^a i) = \psi^i(\omega)$ and that orbit is finite, one value has infinitely many preimages. Compactness supplies a limit point of zeros of $g - c$, so $g \equiv c$. The full proof's work is precisely to construct such a g uniformly on a neighbourhood where rational iteration never meets indeterminacy.

Scope and caveats

The proof depends essentially on Hrushovski's theorem to find the good periodic reduction point; the remaining argument follows the Bell–Ghioca–Tucker interpolation method. It supplies neither an effective bound for N nor a Zariski-dense orbit, and it does not classify the preperiodic points. Its conclusion is instead the robust minimum one can ask for: infinite-order rational dynamics cannot make every algebraic point preperiodic.