

Pulling Back Cohomology Classes and Dynamical Degrees Of Monomial Maps

Jan-Li Lin

arXiv:1010.6285

<http://arxiv.org/abs/1010.6285v1>

One-sentence summary. Lin computes every dynamical degree of a dominant monomial map from the ordered moduli of the eigenvalues of its exponent matrix, by turning toric pullback into complementary minors and then passing to an exterior power.

From an integer matrix to birational dynamics

Let $N \simeq \mathbf{Z}^n$, and let $\psi = (a_{ij}) \in M_n(\mathbf{Z})$. On the dense torus it defines

$$f_\psi(x_1, \dots, x_n) = \left(\prod_{j=1}^n x_j^{a_{1j}}, \dots, \prod_{j=1}^n x_j^{a_{nj}} \right).$$

This extends equivariantly to a rational self-map of any compatible complete toric variety, and is dominant precisely when $\det(\psi) \neq 0$. For a dominant meromorphic self-map of a compact Kähler n -fold, its k -th dynamical degree measures the exponential growth of pullback on $H^{k,k}$:

$$\lambda_k(f) = \lim_{\ell \rightarrow \infty} \|(f^\ell)^*\|^{1/\ell}.$$

Main theorem (Lin, Theorem 1). *Let $f = f_\psi$ be dominant, and order the eigenvalues of ψ , counted with multiplicity, so that $|\mu_1| \geq \dots \geq |\mu_n|$. Then for $1 \leq k \leq n$,*

$$\lambda_k(f) = |\mu_1 \cdots \mu_k|.$$

Thus all higher dynamical degrees, not merely the first, are encoded by the linear action on the cocharacter lattice. The formula is birationally meaningful because dynamical degrees do not depend on the chosen compact model.

Ingredient I: a toric basis that sees minors

Birational invariance permits the convenient model $X = (\mathbf{P}^1)^n$. Its fan has rays $\pm e_i$. For every $\alpha \subset \{1, \dots, n\}$ with $|\alpha| = n - k$, the coordinate orbit closure gives a Chow class, and the corresponding Minkowski weight c_α is 1 on cones contained in $E_\alpha = \text{span}\{e_i : i \in \alpha\}$ and 0 otherwise. These c_α form a basis of $A^k(X)$, dual to complementary coordinate cycles.

The paper's concrete pullback formula (Theorem 4.1) is

$$(f_\psi^*)_{\alpha\beta} = |\det(\psi_{\beta', \alpha'})|, \quad |\alpha| = |\beta| = n - k,$$

where a prime denotes the complementary index set. Apart from absolute values and transpose, this is exactly the matrix of $\bigwedge^k \psi$. It is the bridge from a rational map, whose pullback need not behave functorially under iteration, to ordinary finite-dimensional linear algebra.

Ingredient II: resolve the rational pullback

To establish that formula, refine the fan until f_ψ becomes a morphism and write rational pullback as pullback upstairs followed by toric pushforward. The projection formula reduces an (α, β) entry to intersections of cones lying in $\psi^{-1}E_\beta$ and $E_{\alpha'}$. After a generic translation, the two underlying linear spaces are either disjoint, giving zero, or meet once, giving the product of lattice indices

$$[N : (\psi^{-1}E_\beta \cap N) + N_{\alpha'}] [N : \psi(N) + N_\beta].$$

Lemma 4.3 identifies this product with $|\det(\psi) \det((\psi^{-1})_{\alpha, \beta})|$. Lemma 4.2, the complementary minor identity obtained on exterior powers, converts it to $|\det(\psi_{\beta', \alpha'})|$. This lattice-index calculation is the proof's technical core: it shows that resolving indeterminacy introduces exactly the multiplicities required by the minor, and no others.

Ingredient III: iterate before taking spectral growth

The iterate f^ℓ is the monomial map induced by ψ^ℓ . Applying Theorem 4.1 separately to each iterate gives

$$\lambda_k(f) = \lim_{\ell \rightarrow \infty} \left(\max_{\alpha, \beta} |\det((\psi^\ell)_{\beta', \alpha'})| \right)^{1/\ell} = \lim_{\ell \rightarrow \infty} \|(\bigwedge^k \psi)^\ell\|^{1/\ell}.$$

The last limit is the spectral radius of $\bigwedge^k \psi$. Its eigenvalues are the products $\mu_{i_1} \cdots \mu_{i_k}$, so the largest modulus is $|\mu_1 \cdots \mu_k|$, proving the theorem. Notice the order of operations: one computes the pullback of f^ℓ from ψ^ℓ ; one does not assume the generally false identity $(f^\ell)^* = (f^*)^\ell$ for rational maps.

Diagonal toy model

If $\psi = \text{diag}(a_1, \dots, a_n)$, then $f(x_1, \dots, x_n) = (x_1^{a_1}, \dots, x_n^{a_n})$. Pulling back a coordinate codimension- k cycle indexed by I multiplies its class by $\prod_{i \in I} |a_i|$. After ℓ iterations the factor is its ℓ -th power, and the largest growth comes from the k largest $|a_i|$. The full argument retains this exterior-power picture when eigendirections are not coordinate axes, while the toric resolution accounts for indeterminacy and lattice multiplicities.

Scope and omissions

The theorem requires dominance, equivalently $\det(\psi) \neq 0$, and concerns monomial maps; it is not a formula for arbitrary rational self-maps. The paper also computes all projective degrees of the Cremona involution, but that separate application is omitted here. Likewise, the entropy corollary and the detailed general theory of Minkowski weights are not needed for the central dynamical-degree calculation.