

Rational Curves on K3 Surfaces

Jun Li, Christian Liedtke

arXiv:1012.3777

<http://arxiv.org/abs/1012.3777v2>

One-sentence summary. Li and Liedtke prove that every projective K3 surface of odd Picard rank in characteristic zero contains infinitely many integral rational curves, by combining Picard-rank jumps after reduction with rigidified stable maps that can be transported through moduli.

The odd-rank case of the rational-curve conjecture

An integral rational curve is an irreducible curve whose normalization is $\mathbf{P}^1$. A K3 surface has no moving family of rational curves in characteristic zero, so producing infinitely many such curves requires changing their divisor classes rather than sweeping out the surface.

Main theorem (Li–Liedtke, Theorem 3.3). *Let X be a projective K3 surface over an algebraically closed field of characteristic zero. If the Picard rank $\rho(X)$ is odd, then X contains infinitely many integral rational curves.*

For an ample divisor H , the proof actually produces integral rational curves of arbitrarily large H -degree. This immediately implies infinitude.

Ingredient I: new classes after reduction

By the Bogomolov–Hassett–Tschinkel reduction theorem, it suffices to work over a number field after arranging that every class in $\text{Pic } X$ is defined there. At a density-one set of good places, after a finite field extension, the reductions X_q are ordinary and hence non-supersingular. Theorem 3.1 says that a good non-supersingular reduction in characteristic $p \geq 5$ has even geometric Picard rank. Since the generic Picard rank is odd, $\text{Pic}(X_q)$ contains a class outside $\text{Pic}(X) \otimes \mathbf{Q}$.

Choose such a class effective. Theorem 1.1 represents it by a divisor that is a sum of rational curves, so one component D_q still has class outside the generic Picard space. Proposition 3.2 sharpens this: for every N one can choose q and an integral D_q with

$$[D_q] \notin \text{Pic}(X) \otimes \mathbf{Q}, \quad D_q \cdot H_q \geq N.$$

If degrees stayed below N at infinitely many places, the finite-type scheme of maps $\mathbf{P}^1 \rightarrow X$ of degree $< N$ would supply a characteristic-zero curve specializing to one of them. Its divisor class would lie in the generic Picard group, contradicting the construction. This is the first compactness step in the proof.

Ingredient II: rigidifiers repair reducible curves

A genus-zero stable map is called rigid when its moduli stack is zero-dimensional at that point. On a non-supersingular K3 surface, Lemma 2.6 gives a practical criterion: every non-ghost component maps birationally, ghost components are isolated three-pointed components, and adjacent non-ghost components meet properly. These conditions prevent a one-parameter deformation of the map.

A *rigidifier* is the normalization map $f: \mathbf{P}^1 \rightarrow R$ of an integral nodal rational curve whose class is ample. Theorem 2.9 says that for integral rational curves $D_1, \dots, D_m$ and such an f , some $k \leq m$ makes

$$D_1 + \dots + D_m + kR$$

representable by a rigid stable map. The construction chains copies of the normalization of R through distinct branches at its nodes, then attaches the normalizations of the D_i . Ampleness ensures the necessary intersections, and extra nodes let one reroute attachments when a chosen point is already nodal. Lemma 2.6 then proves rigidity. Proposition 2.8 shows that polarized K3 surfaces admitting rigidifiers in a multiple of the polarization form an open dense locus in moduli.

Ingredient III: deform the special curve back to X

The stable map representing D_q is rigid because X_q is non-supersingular. In a finite étale chart U of the 19-dimensional moduli stack of polarized K3 surfaces, its deformation locus is a divisor B_q : the new line bundle cannot extend over all of U , while Deligne’s formal lifting result makes this divisor flat over $\mathbf{Z}$. Only finitely many primes can yield the same B_q , or else the new class would lift to X . Hence the B_q are infinitely many distinct divisors, and almost all meet the open dense rigidifier locus.

Choose on such a B_q a non-supersingular surface Z carrying both the deformation D of D_q and a rigidifier $R \in |rH|$. For $m \gg 0$, $|mH - D|$ is effective; Theorem 1.1 writes a member as a sum $\sum_i D_i$ of rational curves. Theorem 2.9 adds at most one copy of R per component and produces a rigid stable map of class

$$D + \sum_i D_i + kR = (m + kr)H.$$

The proper relative stable-map stack in this class has dimension at least 19, the dimension of U , and its fibre at this rigid point is zero-dimensional. A component therefore reaches the surface X . After Stein factorization, connectedness of the relevant fibres and rigidity on the non-supersingular reduction keep the special image fixed, so the map on X has an integral component $\tilde{D}$ specializing to a curve that contains D_q . Consequently $\tilde{D} \cdot H \geq D_q \cdot H_q \geq N$.

Why the rigidifier is the hard point

The new curve D_q itself generally cannot lift: its class was chosen precisely outside the generic Picard group. The proof instead moves along its Noether–Lefschetz divisor, completes its class to a polarization multiple, and makes the resulting reducible cycle rigid. That rigidity is what turns properness and the dimension count into controlled transport across moduli. Without the nodal ample rigidifier, multiple or non-reduced components could deform inside the special fibre and the limiting image would no longer be forced to retain D_q .

Scope and caveats

The theorem settles the odd-Picard-rank case in characteristic zero, not the full conjecture for every K3 surface. The paper also proves the analogous statement for non-supersingular odd-rank K3 surfaces in characteristic $p \geq 5$ (Theorem 3.4), with additional ordinary cases in characteristics 2 and 3 noted separately. Its argument is existential: the density-one reductions and moduli transport do not give an effective list or degree bound for the curves on a specified surface.