

Periodic points of birational maps on projective surfaces

Junyi Xie

arXiv:1106.1825

<http://arxiv.org/abs/1106.1825v2>

One-sentence summary. Xie proves that a birational self-map of a projective surface with first dynamical degree greater than one has Zariski-dense non-critical periodic points over every algebraically closed field, by preserving degree growth under specialization and lifting isolated periodic points from finite fields.

Degree growth versus periodic points

Let X be a projective surface over an algebraically closed field, let L be ample, and let $f: X \dashrightarrow X$ be birational. Its first dynamical degree is

$$\lambda_1(f) = \lim_{n \rightarrow \infty} ((f^n)^* L \cdot L)^{1/n} \geq 1.$$

It is independent of L and invariant under birational conjugacy. The qualification “non-critical” excludes periodic points at which the relevant iterate fails to be a local isomorphism, so that indeterminacy and exceptional behavior do not masquerade as honest periodic dynamics.

Main theorem (Xie, Theorem 1.2). *If $f: X \dashrightarrow X$ is birational and $\lambda_1(f) > 1$, then its non-critical periodic points are Zariski dense in X .*

The field may have arbitrary characteristic. For X smooth and the base field of characteristic different from 2 and 3, this is the exponential-growth case of the paper’s complete classification (Theorem 1.1): when $\lambda_1 = 1$, density is instead characterized by finite-order behavior or by an iterate acting trivially on the base of the rational or elliptic fibration dictated by linear or quadratic degree growth.

Ingredient I: make specialization retain $\lambda_1 > 1$

Degrees of iterates can drop in a special fibre, so the inequality $\lambda_1 > 1$ cannot simply be read from the generic map. Xie first lets a birational map act isometrically on the infinite-dimensional hyperbolic space inside the L^2 completion of the Picard–Manin (Riemann–Zariski) space of X . A quantitative local-to-global estimate for a broken geodesic gives an effective criterion: sufficiently large growth from f^n to f^{2n} forces $\lambda_1(f) > 1$. Set

$$q_n = \frac{\deg_L(f^{2n})}{3^{18} \sqrt{2} \deg_L(f^n)}.$$

Corollary 3.2 gives

$$(q_n/2)^{1/n} < \lambda_1(f), \quad \lim_{n \rightarrow \infty} (q_n/2)^{1/n} = \lambda_1(f).$$

Only two finite iterate degrees are therefore needed to certify a prescribed lower bound below $\lambda_1(f)$.

In a projective smooth family, each fixed iterate degree is lower semicontinuous and agrees with the generic degree on a nonempty open set. Applying the finite-iterate criterion simultaneously to f^n and f^{2n} proves Theorem 4.2: the first dynamical degree is lower semicontinuous on the locus of birational fibres. Thus a model of f over a finitely generated ring has closed fibres with $\lambda_1 > 1$.

Ingredient II: periodic points over a finite field

Over the algebraic closure $\overline{\mathbf{F}}_q$, Proposition 6.2 adapts Hrushovski’s Frobenius argument to a birational map ϕ . If the closure of its non-critical periodic points were proper, choose an affine open U outside it on which ϕ is an open embedding. For some $m > 0$, Hrushovski’s intersection estimate makes the graph of ϕ meet the graph of q^m -Frobenius inside $U \times U$, producing

$$\phi(u) = \text{Frob}_q^m(u).$$

Because ϕ is defined over $\mathbf{F}_q$, iteration gives $\phi^\ell(u) = \text{Frob}_q^{m\ell}(u)$. The algebraic point u is defined over a finite extension, hence is periodic, and the choice of U makes it non-critical—a contradiction.

Ingredient III: lift only isolated periodic points

Let a model be flat over a discrete valuation ring. Lemma 6.3 studies $\Delta \cap \Gamma_{\Phi^n}$ in the relative self-product. A smooth non-critical periodic point of the special fibre that lies on no periodic curve belongs to a one-dimensional component of this intersection; that component reaches the generic fibre and yields a non-critical periodic point there. A divisor-avoidance argument shows that these lifts meet every nonempty generic open set.

The exclusion of periodic curves is essential. Cantat’s theorem says that infinitely many periodic curves force the map, after modification, to preserve a fibration. The fibre class is fixed by pullback, and the Hodge index theorem then forces $\lambda_1 = 1$. Hence a map with $\lambda_1 > 1$ has only finitely many periodic curves, exactly the hypothesis needed by the lifting lemma.

Proof roadmap and the hard point

Spread X , L , and f over a regular finitely generated ring. Lower semicontinuity supplies a smooth birational closed fibre with $\lambda_1 > 1$. Proposition 6.2 makes its non-critical periodic points dense; Cantat removes all but finitely many curve components; and Lemma 6.3 lifts a dense supply to the generic fibre. Induction on the transcendence degree over the prime field completes the argument.

The technically hardest step is not the finite-field density but ensuring that exponential degree growth survives reduction. A single iterate can lose degree through new cancellations. The Picard–Manin hyperbolic estimate turns two controlled iterate degrees into a robust lower bound, making the specialization-and-lifting strategy possible.

Scope and omissions

The arbitrary-characteristic theorem above concerns density of non-critical periodic points, not an asymptotic count. The sharper estimate $\#\text{Per}_n(f) \sim \lambda_1(f)^n$, where $\#\text{Per}_n(f)$ counts isolated period- n points with multiplicities, is proved only for nontrivial automorphisms of smooth projective surfaces in characteristic zero (Theorem 1.3). The paper also proves a characteristic-zero dense-orbit theorem and analyzes all three $\lambda_1 = 1$ growth regimes; their separate Saito-formula and fibration arguments are only summarized here.