

# Chow groups of moduli spaces of rank 2 vector bundles on curves with determinant of odd degree

Evgeny Mayanskiy

arXiv:1111.2986

<http://arxiv.org/abs/1111.2986v1>

**One-sentence summary.** Mayanskiy computes the integral Chow groups of the fixed-determinant moduli space of stable rank-two bundles on a curve by transporting the blow-up and projective-bundle formulas through Thaddeus's sequence of flips.

## The moduli problem and the answer

Let  $X$  be a smooth projective curve of genus  $g \geq 2$ . Fix  $d = 2w + 1$  with  $w \geq 2g - 2$  and a line bundle  $\Lambda$  of degree  $d$ . Let  $N$  be the moduli space of rank-two semistable bundles with determinant  $\Lambda$ . Since  $d$  is odd, semistability equals stability. Set

$$n = d - 2g + 2, \quad m = d + g - 1, \quad M_0 = \mathbf{P}^{m-1},$$

and write  $S^k X$  for the  $k$ th symmetric power. All Chow groups below have integer coefficients and  $A^l$  denotes codimension  $l$ .

Define

$$K_l = \bigoplus_{i=0}^{n-2} A^{l+1+i}(N) \oplus \bigoplus_{k=1}^w \bigoplus_{r=0}^{k-2} A^{n-1+l-m+2k-r}(S^k X),$$

$$B_l = A^{n-1+l}(M_0) \oplus \bigoplus_{k=1}^w \bigoplus_{s=0}^{m-2k-2} A^{n-1+l-k-s}(S^k X).$$

**Main theorem** (Mayanskiy). *For every  $l$  there is a natural split exact sequence of abelian groups*

$$0 \longrightarrow K_l \longrightarrow B_l \longrightarrow A^l(N) \longrightarrow 0.$$

*The maps and a splitting are given by standard intersection-theoretic operations and inductively determined polynomials in Chern classes of natural bundles. Consequently every  $A^l(N)$  has a finite natural resolution by direct sums of Chow groups of  $M_0$  and the  $S^k X$ .*

The result is more than an additive generation statement: it records the relations and a section of the quotient map. Its building blocks are Chow groups of a projective space and symmetric powers of the original curve.

## Ingredient I: Thaddeus's birational ladder

Thaddeus constructs a rank- $n$  projective bundle  $M_w \rightarrow N$  and connects it to  $M_0$  by varieties

$$M_{k-1} \xleftarrow{f_k^-} \widetilde{M}_k \xrightarrow{f_k^+} M_k, \quad 1 \leq k \leq w,$$

where both arrows are blow-ups. Their smooth centers are projective bundles over the same base  $S^k X$ :

$$Z_k^- = \mathbf{P}(W_k^-), \quad \mathrm{rk} W_k^- = k, \quad Z_k^+ = \mathbf{P}(W_k^+), \quad \mathrm{rk} W_k^+ = m - 2k.$$

Thus a wall crossing replaces one projective-bundle center by another, while the common exceptional divisor remembers both descriptions.

## Ingredient II: one wall gives generators and relations

For the blow-up  $f: \widetilde{Y} \rightarrow Y$  of a smooth variety along a smooth codimension- $c$  center  $Z$ , with exceptional inclusion  $j$  and projection  $q$ , the standard formula is

$$A^l(\widetilde{Y}) \simeq f^* A^l(Y) \oplus j_* \bigoplus_{a=0}^{c-2} \xi^a q^* A^{l-a-1}(Z).$$

Apply this first to  $f_k^-$  and then to  $f_k^+$ , and expand each  $A^*(Z_k^\pm)$  by the projective-bundle formula over  $S^k X$ . Cancelling the common exceptional terms yields the split exact sequence

$$0 \rightarrow \bigoplus_{r=0}^{k-2} A^{l-m+2k-r}(S^k X) \rightarrow A^l(M_{k-1}) \oplus \bigoplus_{s=0}^{m-2k-2} A^{l-k-s}(S^k X) \rightarrow A^l(M_k) \rightarrow 0. \quad (1)$$

The maps are concrete combinations of pullback, proper pushforward, the center embeddings, and powers of tautological classes. Rewriting powers from the two projective-bundle structures introduces the Chern- and Segre-class polynomials displayed in the paper.

## A first-wall model

At  $k = 1$ , the left sum in (1) is empty because  $Z_1^-$  has rank one over  $X$ . Hence

$$A^l(M_1) \simeq A^l(M_0) \oplus \bigoplus_{s=0}^{m-4} A^{l-1-s}(X).$$

The first wall simply contributes new shifted Chow groups of the curve. At later walls the left side of (1) is nonzero: some classes from  $S^k X$  become relations among the old and new generators. This is the local model for the theorem's two large direct sums.

## Ingredient III: telescope the split sequences

Write (1) schematically as  $0 \rightarrow S_k \rightarrow A^l(M_{k-1}) \oplus R_k \rightarrow A^l(M_k) \rightarrow 0$ . Because every step is split, the intermediate  $A^l(M_k)$  terms can be eliminated without losing exactness or naturality. The  $w$  wall crossings give

$$0 \rightarrow \bigoplus_{k=1}^w S_k \rightarrow A^l(M_0) \oplus \bigoplus_{k=1}^w R_k \rightarrow A^l(M_w) \rightarrow 0. \quad (2)$$

This telescoping is the organizational core of the calculation: each symmetric power occurs at precisely the shifts dictated by the ranks  $k$  and  $m - 2k$  of the two centers.

## Ingredient IV: descend from $M_w$ to $N$

The final space is not  $N$  itself but a rank- $n$  projective bundle over it. The projective-bundle formula, followed by projection to the top power of the tautological class, gives another split sequence

$$0 \rightarrow \bigoplus_{i=0}^{n-2} A^{l+n-1-i}(N) \rightarrow A^{l+n-1}(M_w) \rightarrow A^l(N) \rightarrow 0. \quad (3)$$

Insert (2), with codimension shifted by  $n - 1$ , into (3) and reindex the first sum. The result is exactly  $0 \rightarrow K_l \rightarrow B_l \rightarrow A^l(N) \rightarrow 0$ . Repeatedly resolving the remaining higher-codimension Chow groups of  $N$  terminates and gives the corollary's finite resolution.

## Proof roadmap and the hard point

Start from  $M_0 = \mathbf{P}^{m-1}$ ; at each Thaddeus wall, compute the same exceptional divisor from both blow-up directions; use the two projective-bundle formulas to isolate a split relation (1); telescope all walls to reach  $M_w$ ; and finally use the projective bundle  $M_w \rightarrow N$  to solve for  $A^l(N)$ .

The difficult point is compatibility, not a single vanishing theorem. One must track the two tautological classes on the common exceptional divisor, convert them through Chern and Segre classes of  $W_k^\pm$ , and retain splittings that compose naturally through all  $w$  steps. Without those splittings, the successive short exact sequences would not telescope into the explicit global presentation.

## Scope and omissions

The theorem is an additive description of integral Chow groups, not a presentation of the Chow ring. It reduces the answer to the groups  $A^*(S^k X)$  but does not compute those groups further. The hypotheses  $g \geq 2$ , odd  $d = 2w + 1$ , and  $w \geq 2g - 2$  are those under which the stated Thaddeus ladder and ranks are used. The paper records an inductive formula for the natural morphisms; this note suppresses its full Chern-class coefficients while preserving the source, target, and role of every map.