

Dynamical Degree, Arithmetic Entropy, and Canonical Heights for Dominant Rational Self-Maps of Projective Space

Joseph H. Silverman

arXiv:1111.5664

<http://arxiv.org/abs/1111.5664v2>

One-sentence summary. For a monomial self-map of an algebraic torus, Silverman turns vanishing canonical height into multiplicative character relations, proving that every height-zero point lies in the divisible hull of a proper algebraic subgroup and that arithmetic degrees form a finite, explicitly spectral set.

Three growth invariants

Let $A = (a_{ij}) \in \text{Mat}_N(\mathbf{Z})$ have nonzero determinant and define

$$\varphi_A(x_1, \dots, x_N) = \left(\prod_j x_j^{a_{1j}}, \dots, \prod_j x_j^{a_{Nj}} \right) \quad \text{on } \mathbf{G}_m^N.$$

It extends rationally to $\mathbf{P}^N$. Write $\rho = \rho(A)$ for the spectral radius and let $\ell + 1$ be the largest Jordan-block size among eigenvalues of modulus ρ . Theorem 24 (due independently to Lin and to Jonsson–Wulcan) gives

$$\deg(\varphi_A^n) \asymp n^\ell \rho^n, \quad \delta_{\varphi_A} = \rho.$$

Thus, for $P \in \mathbf{G}_m^N(\overline{\mathbf{Q}})$,

$$\widehat{h}_A(P) = \limsup_{n \rightarrow \infty} \frac{h(\varphi_A^n(P))}{n^\ell \rho^n}, \quad \alpha_A(P) = \limsup_{n \rightarrow \infty} h(\varphi_A^n(P))^{1/n}.$$

Following the paper, if the iterated heights are eventually zero, we set $\alpha_A(P) = 1$ by convention. Proposition 25 proves that $\widehat{h}_A(P)$ is finite, while the general height inequalities give $\widehat{h}_A(P) > 0 \Rightarrow \alpha_A(P) = \delta_{\varphi_A}$.

Call a Jordan subspace *maximal* if its eigenvalue has modulus ρ and its dimension is $\ell + 1$. Let $\bar{r}(A)$ be the number of distinct Galois conjugates of maximal Jordan subspaces. For an algebraic subgroup $G \subset \mathbf{G}_m^N$, let

$$G(\overline{\mathbf{Q}})^{\text{div}} = \{P : P^q \in G(\overline{\mathbf{Q}}) \text{ for some } q \geq 1\}.$$

Main theorem (Silverman, Theorem 27 and Corollaries 29, 32). *Assume $\rho > 1$. There is an algebraic subgroup $G \subset \mathbf{G}_m^N$, depending only on A , with*

$$\dim G \geq N - \bar{r}(A), \quad \{P : \widehat{h}_A(P) = 0\} \subset G(\overline{\mathbf{Q}})^{\text{div}}.$$

Moreover, the orbit of every height-zero point is contained in a proper algebraic subgroup. For every A as above (including $\rho = 1$), if $f_1, \dots, f_s$ are the monic irreducible factors of $\det(T - A)$ and $\rho(f)$ is the largest modulus of a root of f , then

$$\{\alpha_A(P) : P \in \mathbf{G}_m^N(\overline{\mathbf{Q}})\} = \{1, \rho(f_1), \dots, \rho(f_s)\}.$$

This proves the paper's finite-spectrum and dense-orbit conjectures for monomial maps: a Zariski-dense orbit cannot have canonical height zero.

Ingredient I: height zero gives local linear algebra

Choose a number field K containing the coordinates of P and the eigenvalues of A . For a place v , put

$$\log \|P\|_v = {}^t(\log \|x_1\|_v, \dots, \log \|x_N\|_v), \quad \log \|\varphi_A^n(P)\|_v = A^n \log \|P\|_v.$$

The normalized matrices $n^{-\ell} \rho^{-n} A^n$ are bounded. Along a subsequence realizing the limsup, let them converge to a real matrix B . For a real vector u , write $\max^+(u) = \max\{0, u_1, \dots, u_N\}$. If $\widehat{h}_A(P) = 0$, the height decomposition is a sum over v of nonnegative terms $\max^+(B \log \|P\|_v)$, hence each vanishes. The product formula then forces the sharper place-by-place identity

$$B \log \|P\|_v = 0. \tag{1}$$

This is the decisive bridge: a global asymptotic condition becomes an exact linear relation at every place.

Ingredient II: the surviving Jordan directions

On a nonmaximal Jordan block, $n^{-\ell} \rho^{-n} A^n$ tends to zero. On a maximal block V with basis $v_1, \dots, v_{\ell+1}$, its limit kills $W = \langle v_1, \dots, v_\ell \rangle$ but not V/W . Consequently $\ker B = U \otimes_{\overline{\mathbf{Q}}} \mathbf{C}$, where U is obtained by replacing each maximal block by its canonical codimension-one subspace and retaining every other block. This U depends only on A , not on P , the place, or the convergent subsequence.

Ingredient III: Baker descent and characters

Equation (1) initially yields annihilators with algebraic coefficients. Baker's theorem on linear forms in logarithms says that the $\mathbf{Q}$ -space of relations among the logarithms of algebraic numbers is obtained from its $\mathbf{Q}$ -space; in particular it is Galois invariant. Summing the Galois conjugates of $U^\perp$ therefore produces a rational subspace $Y \subset \mathbf{Q}^N$, independent of P and v , with

$$Y \subset (\log\|P\|_v)^\perp \quad \text{for every } v.$$

Let $L = Y \cap \mathbf{Z}^N$. For every $e = (e_1, \dots, e_N) \in L$, all places give

$$\left\| \prod_j x_j^{e_j} \right\|_v = 1.$$

Kronecker's theorem makes this algebraic number a root of unity. Hence some power of P lies in the character kernel

$$G_L = \bigcap_{e \in L} \{X_1^{e_1} \cdots X_N^{e_N} = 1\},$$

which is precisely the asserted divisible-hull containment. The Galois count gives $\dim Y = \bar{r}(A)$ and hence $\dim G_L = N - \bar{r}(A)$.

Toy model and proof roadmap

Take $A = \text{diag}(d, e)$ with integers $d > e \geq 1$. Then $\rho = d$, $\ell = 0$, and the normalized iterates retain only the first coordinate. Thus $\hat{h}_A(x, y) = h(x)$: height zero says that x is a root of unity, while y may be arbitrary. The exceptional set is the divisible hull of the proper subtorus $\{x = 1\}$. This shows why canonical height zero need not mean preperiodic and why the theorem's subgroup, rather than torsion, is the correct conclusion.

In general the proof follows the same pattern: normalize degree growth; extract B ; use height zero and the product formula to obtain (1); read $\ker B$ from maximal Jordan blocks; descend its orthogonal complement using Baker; and turn the resulting integral lattice into a subtorus using Kronecker. Every iterate has height zero and hence lies in $G_L(\overline{\mathbf{Q}})^{\text{div}}$. The orbit lies in the fixed field $K = \mathbf{Q}(P)$; if d is the number of roots of unity in K , then every orbit point Q satisfies $Q^d \in G_L$. Thus the entire orbit lies in the single proper character subgroup G_{dL} .

For the spectrum, first replace P by a coordinatewise power P^d to pass from the finite component group of the smallest orbit-containing algebraic subgroup to its identity component; equation (39) preserves $\alpha_A(P)$. Restriction to this connected invariant subtorus then shows that the arithmetic degree is the spectral radius of a rational invariant subspace, proving one inclusion. Conversely, for an irreducible factor f with $\rho(f) > 1$, write $\det(T - A) = f(T)^e g(T)$ and form the invariant torus $H = g(\varphi_A)(\mathbf{G}_m^N)$. The characteristic polynomial on H is a power of f ; a point with Zariski-dense orbit in H has positive canonical height and arithmetic degree $\rho(f)$. The identity point realizes 1.

Hard point and scope

The hard point is the arithmetic descent from complex limiting directions to integral characters uniform in both P and v ; Jordan theory alone cannot supply it, and Baker's theorem is essential at archimedean places. The paper's canonical height is a limsup, not generally a limit. Under the hypothesis $\rho(A) > 1$, when A is diagonalizable, Corollary 33 strengthens the result to $\hat{h}_A(P) > 0 \iff \alpha_A(P) = \delta_{\varphi_A}$; when $\det(T - A)$ is irreducible over $\mathbf{Q}$ (again with $\rho(A) > 1$), Corollary 31 gives $\hat{h}_A(P) = 0 \iff P$ is preperiodic. Neither stronger equivalence is asserted for an arbitrary matrix.