

Automorphisms of moduli spaces of vector bundles over a curve

Indranil Biswas, Tomás L. Gómez, Vicente Muñoz

arXiv:1202.2961

<http://arxiv.org/abs/1202.2961v1>

One-sentence summary. Biswas, Gómez, and Muñoz recover a curve and the Lie-theoretic data of a general bundle from the intrinsic Hitchin geometry of its fixed-determinant moduli space, forcing every automorphism to come from pullback, tensorization, or dualization.

The moduli space and its automorphisms

Let X be a smooth complex projective curve of genus $g \geq 4$, let $r \geq 2$, and fix a line bundle Λ of degree d . Write $M = M_X(r, \Lambda)$ for the moduli space of stable rank- r bundles E with $\det E \simeq \Lambda$.

Two constructions visibly preserve M . Given $\sigma \in \text{Aut } X$, one may send

$$E \mapsto L \otimes \sigma^* E \quad \text{if} \quad L^r \otimes \sigma^* \Lambda \simeq \Lambda, \quad (1)$$

or

$$E \mapsto L \otimes \sigma^*(E^\vee) \quad \text{if} \quad L^r \otimes \sigma^* \Lambda^{-1} \simeq \Lambda. \quad (2)$$

The degree condition for (2) forces $r \mid 2d$.

Main theorem (Biswas–Gómez–Muñoz, Theorems 1.1 and 5.3). *Every algebraic automorphism of $M_X(r, \Lambda)$ is of type (1) or (2) above.*

The theorem says that the abstract variety M remembers not only its base curve but enough of a general bundle to exclude hidden automorphisms.

Ingredient I: recover the curve from the discriminant

The cotangent bundle T^*M is an open part of the Higgs-bundle moduli space. Its global algebraic functions recover the Hitchin map, up to an automorphism of its affine base,

$$h : T^*M \longrightarrow W = \bigoplus_{k=2}^r H^0(X, K_X^k).$$

The standard fiber dilation induces the weighted $\mathbf{C}^*$ -action on W . Let $D \subset W$ be the locus whose spectral curve is singular. The paper proves intrinsically that $h^{-1}(D)$ is the closure of the union of complete rational curves in T^*M : smooth spectral fibers are open in abelian varieties, whereas a general one-nodal fiber is birationally covered by rational curves. Hence every automorphism of M preserves the Hitchin discriminant.

The largest weight recovers $W_r = H^0(K_X^r)$. For $C = D \cap W_r$, a section s_r lies in C exactly when it has a zero of order at least two. Thus $\mathbf{P}(C)$ is the projective dual of the r -canonical embedding $X \subset \mathbf{P}(W_r^*)$. Duality reconstructs X , giving the Torelli step and the curve automorphism σ induced by an automorphism of M .

Ingredient II: rigidify the Hitchin-base action

After composing away σ , the differential of an automorphism sends T_E^*M to $T_{E'}^*M$ and covers a $\mathbf{C}^*$ -equivariant automorphism f of W . Over a general $s \in W$, this extends across codimension two to an isomorphism of the polarized Prym compactifications of the Hitchin fibers. The polarized Hodge structure then recovers an isomorphism of spectral curves $X_s \simeq X_{f(s)}$ over X .

Inside the total space of K_X , such an isomorphism has fiber coordinate

$$y \mapsto g_1 y + g_2, \quad g_1 \in \mathbf{C}^*, \quad g_2 \in H^0(X, K_X).$$

The spectral equations are trace-free, so their y^{r-1} coefficient s_1 is zero; translation would create such a coefficient. Hence $g_2 = 0$, f is a constant dilation, and after normalizing that dilation one has $f = \text{id}_W$.

Ingredient III: recover a bundle up to duality and twist

For a general E , Proposition 5.1 reconstructs every evaluation kernel

$$\mathcal{E}_x = H^0(\text{End}_0 E \otimes K_X(-x)) \subset T_E^*M$$

from the top Hitchin invariant $h_r = \det$. Surjectivity of evaluation at x follows from $H^0(\text{End}_0 E(x)) = 0$, while the pointwise linear-algebra test

$$\det(A + C) = 0 \text{ for every singular } C \in \text{End}_0(E_x) \implies A = 0$$

characterizes the kernel. Taking the quotient of the resulting subbundle recovers $\text{End}_0 E \otimes K_X$.

The simultaneous vanishing at x of all Hitchin invariants $h_2, \dots, h_r$ recovers the nilpotent cone $\mathcal{N}_{E,x} \subset \text{End}_0(E_x)$. Gerstenhaber’s theorem identifies complete flags in E_x with the maximal linear subspaces of $\mathcal{N}_{E,x}$. Thus the nilpotent-cone bundle determines the complete flag bundle and hence the Lie algebra bundle $\text{End}_0 E$.

Rank two as a toy model

When $r = 2$, the nilpotent cone in $\mathfrak{sl}_2$ is the cone of traceless singular matrices. Its maximal linear subspaces are one-dimensional and are parametrized by lines in a two-dimensional vector space, namely $\text{Fl}(E_x) = \mathbf{P}(E_x)$. This is the smallest instance of the mechanism: invariant polynomials recover a singular matrix cone, and the incidence geometry of that cone recovers flags. Higher rank replaces lines by subspaces of dimension $(r^2 - r)/2$ and uses complete flags.

Ingredient IV and proof roadmap

An isomorphism of the Lie algebra bundles $\text{End}_0 E$ and $\text{End}_0 E'$ has only the inner and dual alternatives for $\mathfrak{sl}_r$. Lifting the associated projective bundles to GL_r therefore gives

$$E' \simeq E \otimes L \quad \text{or} \quad E' \simeq E^\vee \otimes L.$$

The fixed determinant imposes $L' \simeq \mathcal{O}_X$ in the first normalized branch and $L' \simeq \Lambda^2$ in the second. Restoring σ yields exactly (1) and (2), first for general E and then for all E .

In summary: intrinsic rational curves locate the discriminant; its top weight dual recovers X ; Prym–Torelli rigidity kills non-scalar motion of the Hitchin base; determinant evaluation and nilpotent cones recover $\text{End}_0 E$; and Lie theory leaves only tensor or dual-tensor operations.

Hard point and scope

The hardest rigidity step is showing that a priori nonlinear motion of the Hitchin base is only scalar: it passes through compactified generic fibers, polarized Pryms, Torelli for spectral curves, and finally the trace-free condition. The theorem concerns stable bundles on a complex curve with $g \geq 4$; the paper’s Torelli reconstruction alone works for $g \geq 3$. The argument is generic before extending over M , and it relies on standard geometry of Hitchin fibers and on Gerstenhaber’s classification. The paper also explains that the same automorphism group acts on the semistable compactification, but the digest does not reproduce that extension.