

# Uniform bounds for pre-periodic points in families of twists

Alon Levy, Michelle Manes, Bianca Thompson

arXiv:1204.4447

<http://arxiv.org/abs/1204.4447v2>

**One-sentence summary.** For a fixed projective dynamical system over a number field, Levy, Manes, and Thompson bound rational preperiodic points uniformly across all of its twists by controlling the field needed to undo the twist and then applying Northcott's theorem.

## Twists and the uniformity problem

Let  $K$  be a number field and let  $\phi: \mathbf{P}_K^N \rightarrow \mathbf{P}_K^N$  be a morphism of degree  $d \geq 2$ . For  $f \in \mathrm{PGL}_{N+1}(\overline{K})$  write

$$\phi^f = f \circ \phi \circ f^{-1}.$$

A  $K$ -twist of  $\phi$  is a  $K$ -defined map  $\psi$  that is  $\overline{K}$ -conjugate to  $\phi$ , modulo conjugacy by  $\mathrm{PGL}_{N+1}(K)$ . Geometric conjugacy preserves orbit graphs, but it need not preserve which vertices are  $K$ -rational: the conjugating coordinates may live over an extension.

Let

$$A_\phi = \{a \in \mathrm{PGL}_{N+1}(\overline{K}) : \phi^a = \phi\},$$

the finite automorphism group of  $\phi$ .

**Main theorem** (Levy–Manes–Thompson, Theorem 2.8). *There is a constant  $B_\phi$  such that every  $\psi \in \mathrm{Twist}(\phi/K)$  satisfies*

$$\#\mathrm{PrePer}(\psi, \mathbf{P}^N(K)) \leq B_\phi.$$

*The constant is independent of the twist  $\psi$ .*

This is uniformity in one geometric conjugacy class, not the full Morton–Silverman conjecture, whose bound would depend only on  $[K : \mathbf{Q}]$ ,  $N$ , and  $d$ .

## Ingredient I: a twist has bounded descent degree

Choose  $f$  with  $\phi^f = \psi$ , normalize one nonzero matrix entry to 1, and let  $K_f$  be the field generated by the remaining entries. If  $L_f$  is its Galois closure, then for every  $\sigma \in \mathrm{Gal}(L_f/K)$  the  $K$ -rationality of  $\psi$  gives

$$f^{-1}f^\sigma \in A_\phi. \tag{1}$$

Thus Galois conjugates of the coordinate change can differ only by one of the finitely many dynamical symmetries of  $\phi$ .

The map  $\sigma \mapsto f^{-1}f^\sigma$  need not be injective, but each fiber has at most  $[L_f : K_f]$  elements: equality of two images says that the corresponding quotient in the Galois group fixes  $f$ . Counting the source of this map therefore yields the key lemma

$$[K_f : K] \leq \#A_\phi. \tag{2}$$

This is the only place where the twist varies, and (2) removes that variation from the degree of the required field extension.

## Ingredient II: Northcott turns degree into finiteness

Put  $D = \#A_\phi$ . Northcott's theorem says that the preperiodic points of  $\phi$  have bounded height; consequently

$$S_D = \{P \in \mathrm{PrePer}(\phi, \mathbf{P}^N(\overline{K})) : [K(P) : K] \leq D\}$$

is finite. If  $Q \in \mathbf{P}^N(K)$  is preperiodic for a twist  $\psi$ , then  $f^{-1}(Q)$  is preperiodic for  $\phi$  and is defined over  $K_f$ . By (2) it lies in  $S_D$ , and conjugacy is injective on points. Taking  $B_\phi = \#S_D$  proves the theorem.

The proof's economy is the point: Northcott alone is nonuniform in the map, while conjugacy alone loses rationality; bounded descent degree makes the two statements fit exactly.

## A concrete model and the explicit refinement

For  $b \in \mathbf{Q}^*$  consider

$$\phi_b(z) = z + \frac{b}{z}.$$

The change of coordinates  $f_b(z) = \sqrt{b}z$  conjugates  $\phi_1$  to  $\phi_b$ , so every member is a quadratic twist whose conjugating field  $\mathbf{Q}(\sqrt{b})$  has degree at most two. The general theorem already gives an unspecified uniform bound. The paper goes further (Corollary 4.4):

$$\#\mathrm{PrePer}(\phi_b, \mathbf{P}^1(\mathbf{Q})) \in \{2, 4, 6\}.$$

Here  $\infty$  is fixed and  $0 \mapsto \infty$  for every  $b$ ; the representatives  $b = -1$  and  $b = -2$  realize the four- and six-point graphs. The extra argument is genuinely arithmetic. Dynatomic polynomials encode points of exact period. Their parity in  $z$  permits a homogeneous two-variable polynomial, while a coefficient calculation shows that after specialization its leading coefficient is 1 or a prime and its constant coefficient is 1. The rational-root theorem then bounds the possible roots uniformly and rules out exact periods  $n \geq 5$  (Theorem 4.1). Previously established classifications for the two-parameter family  $kz + b/z$  eliminate periods 3, 4 and classify the possible tails, producing the three graphs above. Section 5 treats the distinct twist family

$$\psi_b(z) = -\left(z + \frac{b}{z}\right), \quad b \in \mathbf{Q}^*.$$

Lemma 5.1 gives the analogous extreme-coefficient control at  $k = -1$ ; together with the earlier small-period and tail classifications, it proves that  $\psi_b$  has exactly 2 or 4 rational preperiodic points (Proposition 5.2). The note omits the nearly parallel graph-by-graph proof.

### **Proof roadmap, hard point, and scope**

Undo a twist over  $K_f$ ; encode Galois motion of  $f$  by (1); bound  $[K_f : K]$  by the fixed symmetry group; transport rational preperiodic points back to  $\phi$ ; and apply bounded-height plus bounded-degree finiteness.

The subtle point in the abstract theorem is the descent count: one must use the Galois closure and control the fibers of the cocycle-valued map, since it is not enough merely to observe that  $f^{-1}f^\sigma \in A_\phi$ . The harder computations in the paper concern the sharp value 6, not existence of  $B_\phi$ . In general  $B_\phi$  depends on the fixed pair  $(K, \phi)$  and is not made effective. The explicit classification is only for the stated families over  $\mathbf{Q}$  and uses earlier small-period and tail results; it is not a bound for arbitrary quadratic rational maps.