

On the dynamical and arithmetic degrees of rational self-maps of algebraic varieties

Shu Kawaguchi, Joseph H. Silverman

arXiv:1208.0815

<http://arxiv.org/abs/1208.0815v3>

The problem and the two growth rates

Let K be a number field or a one-dimensional function field of characteristic 0, let X/K be smooth and projective, and let $f: X \dashrightarrow X$ be dominant. Write $X_f(\overline{K})$ for the points whose entire forward orbit avoids the indeterminacy locus. For an ample Weil height h_X , put $h_X^+ = \max\{h_X, 1\}$. The geometric growth rate is

$$\delta_f = \lim_{n \rightarrow \infty} \rho((f^n)^*|_{\text{NS}(X)_{\mathbb{R}}})^{1/n},$$

whereas the upper arithmetic growth of $P \in X_f(\overline{K})$ is

$$\overline{\alpha}_f(P) = \limsup_{n \rightarrow \infty} h_X^+(f^n(P))^{1/n}.$$

The central difficulty is that rational pullback is not functorial: $(f^n)^*$ need not equal $(f^*)^n$.

Main theorem (Theorems 1, 4, and 5). *For every $\varepsilon > 0$ there is $C = C(X, h_X, f, \varepsilon)$ such that, simultaneously for every $P \in X_f(\overline{K})$ and $n \geq 0$,*

$$h_X^+(f^n(P)) \leq C(\delta_f + \varepsilon)^n h_X^+(P).$$

Consequently $\overline{\alpha}_f(P) \leq \delta_f$.

*If moreover f is a morphism and $f^*D \equiv \beta D$ in $\text{NS}(X)_{\mathbb{R}}$ for an $\mathbb{R}$ -divisor D and a real $\beta > \sqrt{\delta_f}$, then*

$$\widehat{h}_{D,f}(P) = \lim_{n \rightarrow \infty} \beta^{-n} h_D(f^n(P))$$

exists, satisfies $\widehat{h}_{D,f} \circ f = \beta \widehat{h}_{D,f}$, and $\widehat{h}_{D,f} = h_D + O(\sqrt{h_X^+})$. If it is nonzero then the lower arithmetic degree is at least β ; hence if $\beta = \delta_f$, the arithmetic degree exists and equals δ_f . When D is ample and K is a number field, $\widehat{h}_{D,f}(P) = 0$ exactly for preperiodic P .

Thus geometric degree growth gives a uniform ceiling for arithmetic orbit growth. The canonical-height assertion is stronger than the classical linearly polarized case: the eigendivisor relation is only algebraic equivalence, and its resulting error is unbounded but of square-root size.

Why the height bound works

- Control non-functorial pullback.** Choose ample divisors $D_1, \dots, D_r$ whose classes form a basis of $\text{NS}(X)_{\mathbb{Q}}$, and let $A(g)$ be the matrix of g^* . Theorem 17 proves

$$\|A(f^{m+n})\| \leq C \|A(f^m)\| \|A(f^n)\|, \quad \|A(f^m)\| \leq C \|A(f)^m\|.$$

The geometric input is the inequality $(g \circ f)^*D \cdot H^{\dim X - 1} \leq f^*(g^*D) \cdot H^{\dim X - 1}$ for nef D, H . After resolving both maps, the discrepancy is effective; positivity on the pseudoeffective cone and equivalence of finite-dimensional norms turn this intersection inequality into the matrix estimate.

- Transfer divisor control to heights.** Proposition 23 gives, for an ample D and a dominant rational g , $h_D(g(Q)) \leq h_{g^*D}(Q) + O(1)$ off the indeterminacy locus. On a resolution $Z \xrightarrow{p} X$, the negativity lemma makes $p^*p_*g^*D - g^*D$ effective and supported over that locus, so its height has the needed lower bound.

- Absorb algebraic-equivalence errors.** If $E \equiv 0$, Lemma 28 supplies the sharp estimate $|h_E(Q)| \ll \sqrt{h_X^+(Q)}$. In the proof, h_X is replaced by the commensurate maximum of the basis heights and normalized so that $h_X \geq 1$. Applying the lemma to $g^*D_k \equiv \sum_i a_{ik}(g)D_i$ then yields, for $Q \notin I_g$,

$$h_X(g(Q)) \leq r \|A(g)\| h_X(Q) + O(\sqrt{h_X^+(Q)}).$$

An elementary recurrence lemma iterates this without letting the square-root term change the exponential rate.

- Choose long blocks.** Set $g = f^{m\ell}$. First choose ℓ so that the spectral radius of $A(f^\ell)$ is at most $(\delta_f + \varepsilon)^\ell$; Jordan normal form contributes only a polynomial in m . Then choose m large enough that all fixed constants and that polynomial have $m\ell$ -th root at most $1 + \varepsilon$. Applying the block estimate to the $m\ell$ residue classes of iterates gives the theorem after a harmless adjustment of ε .

The canonical-height telescope

For the eigendivisor in Theorem 5, Lemma 28 and functoriality for the morphism f give

$$|h_D(f(Q)) - \beta h_D(Q)| \ll \sqrt{h_X^+(Q)}.$$

Consequently, for $N > M$, Theorem 1 bounds the normalized telescoping difference by

$$|\beta^{-N} h_D(f^N(P)) - \beta^{-M} h_D(f^M(P))| \ll \sqrt{h_X^+(P)} \sum_{n>M} \left(\frac{\delta_f + \varepsilon}{\beta^2} \right)^{n/2}.$$

Because $\beta^2 > \delta_f$, one can choose $\varepsilon > 0$ making this a convergent geometric tail. This proves both the Cauchy property and, with $M = 0$, the $O(\sqrt{h_X^+})$ error.

If $\widehat{h}_{D,f}(P) \neq 0$, replace D by $-D$ if necessary and choose an ample H with $H + D$ ample. Then $h_{D+H}(f^n(P)) \geq \beta^n \widehat{h}_{D,f}(P) + o(\beta^n)$, so the lower arithmetic degree is at least β . Finally, when D is ample and the canonical height vanishes, its square-root comparison with h_D bounds the entire orbit height. The orbit lies in the fixed number field $K(P)$, so Northcott finiteness makes it finite.

A rank-one model, the hard point, and scope

If $\text{NS}(X)_{\mathbb{R}} = \mathbb{R}[H]$ and $f^*H \equiv dH$, the proof resembles the scalar recurrence $u_{n+1} \leq du_n + c\sqrt{u_n}$: the error is sublinear, so every $d + \varepsilon$ dominates it exponentially. In higher Picard rank the “coefficient” is instead the non-functorial sequence $A(f^n)$; proving its uniform quasi-submultiplicativity is the hardest new step.

The paper proves an upper bound, not the general existence of $\alpha_f(P)$ nor the conjectural equality $\alpha_f(P) = \delta_f$ for a Zariski-dense orbit. It records several established cases of that conjecture (including morphisms with one-dimensional Néron–Severi space, surface automorphisms, monomial maps for points in the torus, and endomorphisms of abelian varieties), but the general problem remains open in the paper’s scope.