

Cohomology jump loci in the moduli spaces of vector bundles

Botong Wang

arXiv:1210.1487

<http://arxiv.org/abs/1210.1487v1>

Setup: a jump locus inside a singular moduli space

Let X be a connected compact Kähler manifold and let $\mathcal{M}$ be the complex-analytic moduli space of stable holomorphic vector bundles of rank r with vanishing Chern classes. Fix a point represented by E and an integer i , and set $k = \dim_{\mathbb{C}} H^i(X, E)$. The cohomology jump locus

$$\Sigma_k^i(X) = \{F \in \mathcal{M} : \dim_{\mathbb{C}} H^i(X, F) \geq k\}$$

is given its determinantal analytic-subspace structure; this matters because neither $\mathcal{M}$ nor $\Sigma_k^i(X)$ need be reduced. Put $V = H^1(X, \text{End } E)$. Nadel identifies the germ of $\mathcal{M}$ at E with the germ at 0 of the quadratic cone

$$C = \{\eta \in V : \eta \smile \eta = 0 \text{ in } H^2(X, \text{End } E)\},$$

via a canonical local exponential map. Define the linear subspace

$$H = \{\eta \in V : \eta \smile H^{i-1}(X, E) = 0 \text{ and } \eta \smile H^i(X, E) = 0\},$$

where the cup products use $\text{End } E \otimes E \rightarrow E$.

Main theorem (Theorem 1.3). *For sufficiently small neighborhoods U of E in $\mathcal{M}$ and U' of 0 in C , Nadel's exponential isomorphism $U' \simeq U$ restricts to an isomorphism of analytic subspaces*

$$H \cap U' \xrightarrow{\sim} \Sigma_k^i(X) \cap U.$$

Thus, inside the possibly singular quadratic Kuranishi model, the full jump scheme is cut out by linear equations.

This is a higher-rank local analogue of Green–Lazarsfeld's theorem. In rank one the Kuranishi cone is all of $H^1(X, \mathcal{O}_X)$; local linearity under the exponential map implies that irreducible components are translates of subtori. The theorem also gives: a singular point of a rank-one Σ_k^i lies in Σ_{k+1}^i , while for arbitrary rank the germ of Σ_k^i at a point outside Σ_{k+1}^i is quadratic.

The proof mechanism

- 1. Recognize the jump scheme functorially.** Locally represent derived pushforward of a universal family by a complex of finite free sheaves and define Σ_k^i by minors of its adjacent differentials. Proposition 2.1 and Corollary 2.4 show that, for every Artinian local $\mathbb{C}$ -algebra A and every deformation $s: \text{Spec } A \rightarrow \mathcal{M}$ supported at E , the map factors through Σ_k^i exactly when $H^i(X, E_A)$ is a free A -module of rank k . Testing all A , rather than only dual numbers, retains nilpotents and higher infinitesimal structure.
- 2. Write the Kuranishi deformation explicitly.** Stability and vanishing Chern classes give E a flat Hermitian–Einstein metric. Hence V is represented by harmonic $\text{End } E$ -valued $(0, 1)$ -forms, and the Kuranishi family over C has holomorphic structure $\bar{\partial}_E + \tilde{\eta}$. Pulling it back along $\text{Spec } A \rightarrow C$ produces E_A and a well-defined linear space of derivative directions $D(E_A) \subset V$.

3. Convert constant cohomology into two annihilators. Proposition 3.5 proves the decisive equivalence

$$H^i(X, E_A) \text{ free of rank } k \iff D(E_A) \subset H.$$

The proof filters the maximal ideal of A by one-dimensional quotients and uses the resulting exact sequences $0 \rightarrow E \rightarrow E_{j+1} \rightarrow E_j \rightarrow 0$. Their boundary maps are cup products with the corresponding derivative classes. Vanishing in degree i prevents classes of $H^i(X, E)$ from being killed; vanishing from degree $i - 1$ prevents new boundaries from entering H^i .

The analytic input making the induction exact is Lemma 3.6: contracting a harmonic $\text{End } E$ -valued $(0, 1)$ -form with a harmonic E -valued $(0, j)$ -form is again harmonic. Thus a zero cup-product class is already represented by the zero harmonic form, so the expected lifted cohomology classes remain $\bar{\partial}_A$ -closed at every Artinian order.

4. Identify the germ. Steps 1–3 say that $\Sigma_k^i \cap U$ and $H \cap U'$ select exactly the same A -points for every Artinian local A . Their completed local rings, and hence their analytic germs under the exponential map, coincide. Shrinking U, U' gives the asserted isomorphism.

Dual-number model and limitations

For $A = \mathbb{C}[\epsilon]/(\epsilon^2)$, a tangent direction $\eta \in V$ deforms $\bar{\partial}_E$ to $\bar{\partial}_E + \epsilon\eta$. A harmonic class $[\xi] \in H^i(X, E)$ lifts only if $\eta \smile [\xi] = 0$; simultaneously, $\eta \smile H^{i-1}(X, E) = 0$ ensures that no new boundary enters degree i . This first-order picture explains the two annihilators. The hard point of the paper is proving that the same linear condition controls every Artinian thickening, hence the nonreduced germ rather than merely its tangent space.

The result is local, assumes stability and vanishing Chern classes on a compact Kähler manifold, and concerns points with $\dim H^i(X, E) = k$. It does not claim that higher-rank jump loci are globally translates of subtori; the ambient quadratic cone may remain singular.