

Curves and cycles on K3 surfaces

Daniel Huybrechts (with an appendix by Claire Voisin)

arXiv:1303.4564

<http://arxiv.org/abs/1303.4564v3>

From rational curves to constant cycle curves

Let X be a projective K3 surface over an algebraically closed field and let $c_X \in \mathrm{CH}_0(X)$ be the Beauville–Voisin class of degree one. For an integral curve $C \subset X$, view its generic point as a closed point of $X_{k(C)}$ and set

$$\kappa_C = [\eta_C] - (c_X)_{k(C)} \in \mathrm{CH}_0(X_{k(C)}).$$

The curve is a *constant cycle curve* if κ_C is torsion; its order is the exact order of this class. Over an uncountable field this is equivalent to every closed point of C representing c_X in $\mathrm{CH}_0(X)$. Rational curves have order one, but the paper constructs non-rational examples, even of order one.

Main theorem (Huybrechts, Proposition 5.1). *Let X be a projective K3 surface over an algebraically closed field of characteristic zero. For every line bundle L and positive integer n , the linear system $|L|$ contains at most finitely many constant cycle curves of order n . Consequently the same holds with “order n ” replaced by “order at most n ”.*

This parallels finiteness of rational curves in a fixed linear system, but the order bound is essential: elliptic-fibration constructions produce dense unions of constant cycle curves whose orders are unbounded.

Ingredient I: a cycle with an Abel–Jacobi invariant

Let $f_C: \tilde{C} \rightarrow C \hookrightarrow X$ be the normalization followed by inclusion. Choose a degree-one divisor $\sum n_i y_i$ on $\tilde{C}$ and a point x_0 representing c_X , and form the homologically trivial cycle

$$Z_C = \Delta_{\tilde{C} \rightarrow X} - C \times \left(\sum n_i y_i \right) - \{x_0\} \times \tilde{C} \quad \text{on } X \times \tilde{C}.$$

Its restriction to the generic fiber is κ_C (Lemma 5.2). Project the Abel–Jacobi class of Z_C to the intermediate Jacobian built from the transcendental lattice $T(X) = \mathrm{NS}(X)^\perp$; denote the result by $\mathrm{AJ}_{\mathrm{tr}}(\kappa_C)$. Lemma 5.4 shows that this projection is independent of the auxiliary degree-one divisor. Corollary 5.5 supplies the necessary implication

$$n\kappa_C = 0 \quad \implies \quad n \mathrm{AJ}_{\mathrm{tr}}(\kappa_C) = 0. \quad (1)$$

Thus constant cycle curves of order dividing n lie in a torsion-zero locus that Hodge theory can detect.

Ingredient II: globalize over a family

Suppose a positive-dimensional locally closed $T \subset |L|$ parametrizes integral curves of one analytic singularity type. After a generically finite base change, the family admits a simultaneous normalization $p: \tilde{C} \rightarrow T_0$ over a dense open subset and an evaluation map $f: \tilde{C} \rightarrow X$, whose fiber is f_C . Set

$$\pi = p \circ \mathrm{pr}_2: X \times \tilde{C} \longrightarrow T_0.$$

If $L^2 = 2g - 2$ and $g > 1$, choose the auxiliary degree-one line bundle L_0 fiberwise so that

$$L_0^{2g-2} \simeq f_C^* L.$$

Lemma 5.6 then kills the algebraic part of the Abel–Jacobi class; the genus-one case is handled by the paper’s indicated adaptation. Write $\mathcal{Z}$ for the global cycle on $X \times \tilde{C}$ whose fiber is Z_{C_t} , and $\mathcal{J}^3 \rightarrow T_0$ for the family of intermediate Jacobians of π . The fiberwise Abel–Jacobi classes define a normal function

$$\nu \in \Gamma(T_0, \mathcal{J}^3).$$

Write ν_{tr} for its projection to the transcendental Jacobian family. The natural map from the full intermediate Jacobian to its transcendental and algebraic parts is a finite isogeny of degree N . Since the algebraic part is zero, the relevant zero loci satisfy

$$Z(n\nu_{\mathrm{tr}}) \subset Z(nN\nu). \quad (2)$$

Ingredient III: the boundary class cannot vanish

The decisive Lemma 5.8 says that the connecting class

$$\delta(\nu) \in H^1(T_0, R^3 \pi_* \mathbf{Z})$$

is non-torsion. After restricting to a curve in T , the global class $[\mathcal{Z}]$ is nonzero in $[C]^\perp \otimes H^2(\tilde{\mathcal{C}}_{T_0}, \mathbf{Z})$. Indeed, $(2g-2)[\mathcal{Z}]$ is the pullback of

$$(2g-2)[\Delta_X] - [C] \times [C],$$

whose action on $[C]^\perp$ is $(2g-2)$ id; dominance of the normalized total family keeps the transcendental part nonzero. Yet its restriction to each fiber is zero because each Z_{C_t} is homologically trivial. The Leray spectral sequence therefore sends this global class to a nonzero class in $H^1(T_0, R^3\pi_*\mathbf{Z})$, and the same argument for every multiple proves non-torsion. This Leray image is precisely $\delta(\nu)$.

Ingredient IV: algebraicity forces finiteness

By the Brosnan–Pearlstein and Saito theorem, the zero locus of the admissible normal function $nN\nu$ is algebraic. If it had a positive-dimensional component, repeating the construction over that component would make $nN\delta(\nu)$ vanish, contradicting the preceding non-torsion statement. Hence its zero locus is zero-dimensional and finite. In view of (1)–(2), only finitely many order- n constant cycle curves occur on each such family. Finally one stratifies $|L|$ into finitely many pieces by singularity type and number of integral components, which proves the theorem. The Picard-rank-one assumption used to simplify the displayed decomposition is removed as explained in Remark 5.10.

Toy model, hard point, and scope

For a rational curve C , the generic-point class already comes from $\mathrm{CH}_0(X)$ because $\mathrm{CH}_0(X \times \mathbf{P}^1)$ splits by the projective bundle formula; hence $\kappa_C = 0$. At the opposite extreme, on an elliptic K3 surface the multisection swept out by fiberwise m -torsion points has order dividing m . These examples isolate exactly why a bounded-order hypothesis is needed.

The hard step is not constructing the normal function but proving that its topological boundary remains non-torsion on every positive-dimensional subfamily; that is what turns algebraicity into finiteness. Characteristic zero is essential to this proof and to the statement as written: unirational K3 surfaces over $\overline{\mathbf{F}}_p$ can have infinitely many rational, hence order-one constant cycle, curves in one linear system. The paper also studies fixed curves of non-symplectic automorphisms, finite fields, and a genus-201 example; these illustrate the theory but are not needed for Proposition 5.1.