

Moduli spaces of quadratic rational maps with a marked periodic point of small order

J. Blanc, J. K. Canci, N. D. Elkies

arXiv:1305.1054

<http://arxiv.org/abs/1305.1054v2>

Marked cycles as moduli surfaces

Let Rat_2 be the affine variety of degree-two endomorphisms of $\mathbf{P}^1$. For $n \geq 1$, let $\widetilde{\mathcal{M}}_2(n)$ parametrize pairs consisting of $f \in \text{Rat}_2$ and an ordered cycle $(p_1, \dots, p_n)$ of *exact* length n . Quotienting simultaneous conjugation by PGL_2 gives the surface $\mathcal{M}_2(n)$. The automorphism σ_n shifts the marked point around the cycle; thus $\mathcal{M}_2(n)/\langle \sigma_n \rangle$ forgets which point of the orbit was marked.

Main theorem (Blanc–Canci–Elkies, Theorem 1). *For $1 \leq n \leq 5$, both $\mathcal{M}_2(n)$ and $\mathcal{M}_2(n)/\langle \sigma_n \rangle$ are rational over $\mathbf{Q}$. The surface $\mathcal{M}_2(6)$ is smooth and affine and is birational to a projective surface of general type, while $\mathcal{M}_2(6)/\langle \sigma_6 \rangle$ is rational over $\mathbf{Q}$. Nevertheless,*

$$\mathcal{M}_2(6)(\mathbf{Q})$$

is infinite.

The change at period six is therefore geometric rather than an obstruction to rational six-cycles: the marked surface becomes of general type even though it contains explicit curves with infinitely many rational points.

Ingredient I: normalize the orbit

For $n \geq 3$, conjugation places the first three orbit points at $0, \infty, 1$. Writing the remaining points as $x_1, \dots, x_{n-3}$, the first five orbit equations determine the six homogeneous coefficients of f up to scale. Lemma 2.1 proves that for $n \geq 5$ the projection

$$\mathcal{M}_2(n) \longrightarrow \mathbf{A}^{n-3}, \quad (f, p_1, \dots, p_n) \longmapsto (x_1, \dots, x_{n-3}),$$

is an isomorphism onto a locally closed affine surface. Distinctness of the orbit points and nonvanishing of the resultant are the open conditions; the remaining orbit equations are closed conditions.

For $n = 5$ there are no remaining equations, so the image is an open subset of $\mathbf{A}^2$. The cases $n \leq 4$ are handled by similar explicit normal forms. Rationality of the cyclic quotients is not automatic over $\mathbf{Q}$: the paper resolves the shift action and computes invariant function fields (for $n = 5$ via the degree-five del Pezzo surface obtained by blowing up four points of $\mathbf{P}^2$).

Ingredient II: the period-six compactification

For $n = 6$ the affine-coordinate equation is unwieldy. A birational change of variables instead realizes $\mathcal{M}_2(6)$ as an open subset of the quintic surface $S_6 \subset \mathbf{P}^3$ with

$$W^2 F_3(X, Y, Z) = F_5(X, Y, Z),$$

where

$$\begin{aligned} F_3 &= (X + Y + Z)^3 + X^2 Z + XY^2 + YZ^2 + 2XYZ, \\ F_5 &= Z^3 X^2 + X^3 Y^2 + Y^3 Z^2 - XYZ(YZ + XY + XZ). \end{aligned}$$

Lemma 4.1 identifies the boundary exactly: nine lines and fourteen conics, equivalently the locus where the coordinate conditions needed for six distinct points or a nonzero resultant fail. It also makes the shift a linear cyclic permutation of X, Y, Z , accompanied by a sign on W . Consequently all singularities of S_6 lie on the boundary, proving that the moduli open set is smooth; the boundary is cut out by a homogeneous polynomial, proving affineness.

Ingredient III: why the surface is of general type

The quintic has eleven isolated singularities: one triple point and ten double points. Blow them up in $\mathbf{P}^3$, with H the pullback of a hyperplane and E_1 lying over the triple point. For the smooth strict transform $\widehat{S}_6$, adjunction gives

$$K_{\widehat{S}_6} = (H - E_1)|_{\widehat{S}_6}.$$

This linear system is the projection $[W : X : Y : Z] \dashrightarrow [X : Y : Z]$ and has two-dimensional image. Hence $\widehat{S}_6$, and therefore S_6 , is of general type. The same double cover explains the quotient: the cube of the shift is the deck involution, and the residual order-three cyclic action on $\mathbf{P}^2$ has rational invariant field. Thus the full orbit-unmarked quotient is rational even though the covering surface is of general type.

Ingredient IV: infinitely many rational points

On the hyperplane $Z = 0$, the equation becomes

$$W^2(X^3 + 3X^2Y + 4XY^2 + Y^3) = Y^2X^3.$$

This curve is birational to

$$E : y^2 = x^3 + 4x^2 + 3x + 1.$$

The point $P = (0, 1)$ has infinite order: the duplication formula gives $2P = (-7/4, 13/8)$, whose nonintegral coordinates exclude torsion by Nagell–Lutz. Thus $E(\mathbf{Q})$ is infinite. Since E meets the twenty-three boundary curves only finitely often, infinitely many of its rational points lie in $\mathcal{M}_2(6)$. Cyclic symmetry supplies three such elliptic curves. A separate rational curve in S_6 gives an explicit one-parameter family whose maps have both a rational six-cycle and a rational fixed point.

Roadmap, hard point, and scope

Normalize a marked orbit; eliminate the map coefficients; replace the bad affine closure at period six by the explicit double-cover quintic; identify the full boundary; compute the canonical system after resolving the eleven singularities; then exhibit a positive-rank elliptic curve that is not contained in the boundary.

The hard point is the explicit birational model together with its boundary: without the boundary calculation, singular points or degenerate maps could be mistaken for genuine moduli points. General type does not prove the Bombieri–Lang prediction that rational points are non-dense, and the paper does not classify all rational points. It does prove finiteness of S -integral points for a fixed number field and finite set of places, a statement compatible with the infinite rational curves above.