

# Fano varieties with small non-klt locus

Mauro C. Beltrametti, Andreas H3ring, Carla Novelli

arXiv:1309.5342

<http://arxiv.org/abs/1309.5342v2>

---

## Index versus the non-klt locus

A Fano variety here is a normal projective complex variety  $X$  with  $-K_X$  Cartier and ample. Its index is the largest  $k$  for which

$$-K_X \simeq kH$$

with  $H$  a Cartier divisor. Write  $\text{Nklt}(X)$  for the locus where  $X$  is not Kawamata log terminal.

**Main theorem** (Beltrametti–H3ring–Novelli, Theorems 1.3 and 1.4). *If  $\text{Nklt}(X) \neq \emptyset$ , then*

$$\dim \text{Nklt}(X) \geq k - 1. \tag{1}$$

*Equality holds precisely when*

$$(\text{Nklt}(X), \mathcal{O}_{\text{Nklt}(X)}(H)) \simeq (\mathbf{P}^{k-1}, \mathcal{O}_{\mathbf{P}^{k-1}}(1)). \tag{2}$$

*In this case  $X$  is log canonical and is a generalized cone with this linear space as vertex. Moreover, if  $X$  is log canonical and  $\dim \text{Nklt}(X) = k$ , then the polarized non-klt locus is either  $(\mathbf{P}^k, \mathcal{O}(1))$  or  $(Q^k, \mathcal{O}_{Q^k}(1))$ , where  $Q^k \subset \mathbf{P}^{k+1}$  is a possibly reducible hyperquadric.*

The point is that no Cohen–Macaulay hypothesis is imposed. This matters, for instance, for cones over abelian varieties.

## Ingredient I: expose the bad discrepancies

Take the canonical modification  $\mu : X' \rightarrow X$ . It has canonical singularities and

$$K_{X'} = \mu^* K_X - E, \quad E = - \sum a_i E_i \geq 0, \quad a_i \leq -1, \tag{3}$$

where  $E$  maps onto  $\text{Nklt}(X)$ ; because  $K_X$  is Cartier, the non-Gorenstein locus of  $X'$  lies in  $E$ . Lemma 1.12 produces a  $K_{X'}$ -negative extremal ray  $R$  with  $E \cdot R > 0$  and  $\mu^* H \cdot R > 0$ . Let  $\varphi : X' \rightarrow Y$  be its contraction and let  $F$  be a component of a positive-dimensional fibre.

The restriction of  $\mu$  to  $F \cap E$  is finite. Hence

$$\dim \text{Nklt}(X) \geq \dim(F \cap E) \geq \dim F - 1. \tag{4}$$

On the other hand, from  $-K_{X'} \sim_{\mathbf{Q}} k\mu^* H + E$ , the nefvalue  $r = r(\varphi, \mu^* H)$  satisfies  $r > k$ . The fibre-dimension estimate in adjunction theory gives  $\dim F \geq r - 1 > k - 1$ . Combining this with (4) and integrality of dimension proves (1).

## **Ingredient II: equality forces a cone**

Assume  $\dim \mathrm{Nklt}(X) = k - 1$ . If  $\varphi$  were birational, the sharper fibre estimate would give  $\dim F \geq \lfloor r \rfloor \geq k$ , so equality would hold throughout (4) and  $F \not\subset E$ . Lemma 2.1, proved after localizing on  $Y$  by cutting fibres with general relative hyperplane sections of  $\mu^*H$ , then forces  $r = \lfloor r \rfloor$ . The equality dimensions force  $r = k$ , contradicting  $r > k$ . Thus  $\varphi$  is of fibre type.

Adjunction theory now identifies  $\varphi$  as a  $\mathbf{P}^k$ -bundle. On a general fibre  $F \simeq \mathbf{P}^k$ , adjunction gives

$$\mathcal{O}_F\left(\sum a_i E_i\right) \simeq \mathcal{O}_{\mathbf{P}^k}(-1). \quad (5)$$

Since the summands are anti-effective Cartier divisors near  $F$ , exactly one exceptional component has coefficient  $-1$  and meets  $F$  in a hyperplane; every other component must vanish, or its image would make the non-klt locus too large. Thus  $K_{X'} = \mu^*K_X - E_1$ , the pair  $(X', E_1)$  is log canonical, and so is  $X$ . The two maps

$$X \xleftarrow{\mu} X' \xrightarrow{\varphi} Y$$

give the generalized-cone structure: the relative hyperplane  $E_1$  is contracted to the linearly polarized vertex (2).

## **Ingredient III: the next-dimensional locus**

Now suppose  $X$  is log canonical and  $\dim \mathrm{Nklt}(X) = k$ . Subadjunction first shows that every lc centre has dimension at least  $k - 1$ , with equality only for a linearly polarized  $\mathbf{P}^{k-1}$ . For a minimal  $k$ -dimensional centre  $W$ , Kawamata subadjunction makes  $(W, \Delta_W)$  a log Fano pair of dimension and index  $k$ ; the high-index classification makes  $W$  either  $\mathbf{P}^k$  or an integral quadric.

If  $W$  is not minimal, the weak subadjunction Lemma 3.1 puts the inverse image of its codimension-one lc centre inside the boundary on the normalization  $W^n$ . The same classification gives  $W^n \simeq \mathbf{P}^k$  and identifies that boundary with one hyperplane. The induced degree-one map on the hyperplane, together with semi-normality, proves that  $W$  itself is normal. Finally Nadel vanishing makes  $\mathrm{Nklt}(X)$  connected. If it is reducible, there are exactly two  $\mathbf{P}^k$  components meeting along their unique  $\mathbf{P}^{k-1}$  lc centre: this is the reducible quadric.

## **Toy model, hard point, and scope**

Let  $Y$  have trivial canonical bundle and let  $L$  be very ample. On  $X' = \mathbf{P}(\mathcal{O}_Y^{\oplus k} \oplus L)$  the tautological map contracts  $E \simeq Y \times \mathbf{P}^{k-1}$  to a linear  $\mathbf{P}^{k-1}$ . Its image is an lc Fano variety of index  $k$  and realizes the equality case: the projective-bundle fibres become the rulings of the cone and  $E$  their relative hyperplane.

The technical difficulty is extracting this rigid contraction before log canonicity is known; the canonical modification and the integrality lemma for  $r$  bridge that gap. The paper goes further under additional hypotheses, classifying general fibres of maximal rationally connected fibrations and constructing examples. It does not assert that every  $\dim \mathrm{Nklt}(X) = k$  case has the same global fibre-space description without the stated MRC-dimension or ladder assumptions.