

Variation of the canonical height in a family of rational maps

Dragos Ghioca, Niki Myrto Mavraki

arXiv:1309.5682

<http://arxiv.org/abs/1309.5682v1>

A height theorem across a bad fibre

Fix $d \geq 2$ and consider the family

$$f_t(z) = \frac{z^d + t}{z} = z^{d-1} + \frac{t}{z} \quad \text{over } \overline{\mathbf{Q}}(t).$$

For a rational section $c \in \overline{\mathbf{Q}}(t)$, write $\widehat{h}_f(c)$ for its canonical height on the generic fibre and $\widehat{h}_{f_\lambda}(c(\lambda))$ for the specialized height. The fibre at $\lambda = 0$ is genuinely bad: its degree is $d - 1$, not the generic degree d .

Main theorem (Ghioca–Mavraki, Theorem 1.1). *For every $c \in \overline{\mathbf{Q}}(t)$,*

$$\widehat{h}_{f_\lambda}(c(\lambda)) = \widehat{h}_f(c) h(\lambda) + O_c(1) \quad (\lambda \in \overline{\mathbf{Q}}),$$

where the bound is independent of λ . Equivalently, the left-hand side is a Weil height on $\mathbf{P}^1$ associated to the $\mathbf{Q}$ -divisor $\widehat{h}_f(c) \cdot \{\infty\}$.

Call–Silverman gives an $o(h(\lambda))$ error in general families. Here the error is bounded although the resultant vanishes at $t = 0$; the proof must therefore control every v -adic approach to this singular parameter.

Ingredient I: homogeneous iterates and generic degree

Choose a number field K containing the coefficients of c , and write $c = A/B$ with coprime $A, B \in K[t]$. After a special cancellation in the first iterate when $A(0) = 0$, define coprime pairs

$$A_{n+1} = A_n^d + tB_n^d, \quad B_{n+1} = A_nB_n^{d-1}, \quad f_t^n(c) = [A_n : B_n]. \quad (1)$$

Propositions 3.1 and 3.3 prove, in both cases $c(0) \neq 0$ and $c(0) = 0$, that $\deg A_n > \deg B_n$ for $n \geq 1$ and that no common factor develops. Consequently

$$\widehat{h}_f(c) = \frac{\deg_t(f_t(c))}{d} = \frac{\deg_t(f_t^2(c))}{d^2}. \quad (2)$$

This exact degree identity later converts an ordinary parameter height into the main term of the theorem.

Ingredient II: freeze the escape rate after two steps

For a place v put

$$M_{n,v}(\lambda) = \max\{|A_n(\lambda)|_v, |B_n(\lambda)|_v\}, \quad G_{v,\lambda}(c) = \lim_{n \rightarrow \infty} d^{-n} \log M_{n,v}(\lambda).$$

The elementary telescoping lemma (Proposition 5.1) says that uniform bounds on $M_{n+1,v}/M_{n,v}^d$ control the difference between this limit and any finite normalized iterate. The central local assertion, Proposition 4.5, is the stronger uniform estimate

$$|G_{v,\lambda}(c) - d^{-2} \log M_{2,v}(\lambda)| \leq C_{v,c}, \quad (3)$$

valid for $\lambda \neq 0$ and $c(\lambda) \neq 0, \infty$, with $C_{v,c}$ independent of λ . At all but finitely many non-archimedean places, good reduction makes the normalized norm constant already after the first iterate, so the corresponding error in (3) is zero.

Ingredient III: three local regimes

The hard work proves (3). First the authors treat a constant starting point α . Upper and lower recurrences for $M_{n,v}$ are combined with orbit estimates, separately for $d > 2$ and $d = 2$; in degree two, infinity is not superattracting, so the large-parameter argument needs a longer initial orbit segment.

For nonconstant c , set $\alpha = f_\lambda(c(\lambda))$ and use the scaling identity that relates its homogeneous iterates to those of $c(\lambda)$. Proposition 6.5 splits the parameter line into

$$|\lambda|_v > Q, \quad R \leq |\lambda|_v \leq Q, \quad 0 < |\lambda|_v < R. \quad (4)$$

Leading terms control the first range, the general telescoping bound controls the compact middle range, and the small range is divided according as $c(0)$ is finite nonzero, infinity, or zero (with $c'(0)$ zero or nonzero). In every case either $c(\lambda)$ or its first image enters a constant-point estimate with constants depending only on v, A, B . This case analysis is precisely what prevents the bad fibre from destroying uniformity.

Global assembly and an effective special case

Sum (3) over embeddings and places. Only a finite set S , determined by c , contributes an error. The height machine applied to the rational map $t \mapsto [A_2(t) : B_2(t)]$, whose degree is $d^2 \widehat{h}_f(c)$ by (2), gives

$$d^{-2} h([A_2(\lambda) : B_2(\lambda)]) = \widehat{h}_f(c) h(\lambda) + O_c(1),$$

and the theorem follows.

For the toy case $c(t) = \alpha \neq 0$, one has $\widehat{h}_f(\alpha) = 1/d$. If ℓ counts the non-archimedean places of $\mathbf{Q}(\alpha)$ where $|\alpha|_v \notin \{0, 1\}$, Theorem 1.2 makes the bound explicit:

$$\left| \widehat{h}_{f_\lambda}(\alpha) - \frac{h(\lambda)}{d} \right| < 3d(1 + \ell + 2h(\alpha)).$$

Thus a preperiodic specialization of a root of unity must satisfy $h(\lambda) < 3d^2$. The paper does not classify such parameters, nor does it claim the same bounded variation for arbitrary rational-map families; its most delicate estimates are tailored to the unique bad fibre and to the form of (1).