

Preperiodic points for rational functions defined over a global field in terms of good reduction

Jung Kyu Canci, Laura Paladino

arXiv:1403.2293

<http://arxiv.org/abs/1403.2293v5>

Uniformity from good reduction

Let K be a global field, S a nonempty finite set of places containing all archimedean ones, and $\phi : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ an endomorphism with good reduction outside S . Put $s = |S|$, let $p = \text{char } K$, and set

$$D = [K : \mathbf{Q}] \quad (p = 0), \quad D = [K : \mathbf{F}_p(t)] \quad (p > 0).$$

For $P \in \mathbf{P}^1(K)$, write $\mathcal{O}_\phi(P) = \{\phi^n(P) : n \geq 0\}$.

Main theorem (Canci–Paladino, Theorem 1). *If P is preperiodic, then $|\mathcal{O}_\phi(P)| \leq \eta(p, D, s)$, where*

$$\begin{aligned} \eta(0, D, s) &= \max\{(2^{16s-8} + 3)[12s \log(5s)]^D, [12(s+2) \log(5s+5)]^{4D}\}, \\ \eta(p, D, s) &= (ps)^{4D} \max\{(ps)^{2D}, p^{4s-2}\} \quad (p > 0). \end{aligned} \tag{1}$$

In particular, this single-orbit bound is independent of $\deg \phi$.

Good reduction means that a homogeneous presentation $[F : G]$ over the local ring retains its degree after reduction, equivalently its resultant is a unit. The theorem is a good-reduction form of uniform boundedness valid in both number-field and function-field settings.

Ingredient I: distances and radical coordinates

For a nonarchimedean place $\mathfrak{p}$, the logarithmic chordal distance

$$\delta_{\mathfrak{p}}([x_1 : y_1], [x_2 : y_2]) = v_{\mathfrak{p}}(x_1 y_2 - x_2 y_1) - \sum_{i=1}^2 \min\{v_{\mathfrak{p}}(x_i), v_{\mathfrak{p}}(y_i)\} \tag{2}$$

is nondecreasing under a map of good reduction. On a cycle it is translation invariant, and points whose index difference is prime to the cycle length have the same distance as two consecutive points (Propositions 2.1–2.3).

The ring of S -integers need not be principal. Section 2.3 passes to a finite extension L/K and radical groups $\sqrt{R_S}$ and $\sqrt{R_S^*}$ so that every K -point has coprime integral coordinates, without inserting the class number into the final constants. In these coordinates, equalities of distances become literal divisibility relations and hence equations in a finite-rank unit group.

Ingredient II: first bound the cycle

For a function field, Lemma 3.1 finds a good place with residue field of size at most $(ps)^{2D} - 1$. Morton–Silverman–Zieve reduction theory then writes the period as $n = ap^e$ after replacing ϕ by an iterate, with $a \leq (ps)^{4D} - 1$. After conjugating one cycle point to $[0 : 1]$, write

$$P_i = [A_i x_1 : y_i], \quad A_i = 1 \quad \text{if } (i, p) = 1. \quad (3)$$

If all relevant A_i are units, pairwise distinct reductions give $p^e \leq (ps)^{2D}$. Otherwise the first nonunit A_{p^α} converts many points of the cycle into solutions of

$$A_{p^\alpha} X - Y = 1. \quad (4)$$

The equation is not S -trivial, so Theorem 4 bounds its solutions even in positive characteristic, where Frobenius can otherwise generate infinitely many S -unit solutions. Counting the indices in (4) bounds p^e and proves Theorem 7. For number fields, the analogous cycle bound is the earlier Morton–Silverman estimate, and a finite-rank two-variable unit theorem replaces Theorem 4.

Ingredient III: control tails in each characteristic

First let K be a function field. Choose the small good residue field from Lemma 3.1 and let P_0 be the first periodic point in the orbit. If P_0 is indifferent, Lemma 4.3 makes all strictly preperiodic points in its tail distinct after reduction, so the residue-field size bounds the tail.

If P_0 is attracting, its cycle has at most $(ps)^{2D}$ points. Pass to that bounded iterate, normalize $P_0 = [0 : 1]$, and write the tail as

$$P_{-m+1} \mapsto \cdots \mapsto P_{-1} \mapsto P_0, \quad (5)$$

Lemma 4.1 orders all distances to P_0 . A block on which the distance is constant has at most $(ps)^{2D}$ points. At the first strict valuation drop, coprime coordinates give, for every later index b , an equation

$$\frac{x_1}{x_a y_1 - x_1 y_a} v_b - \frac{x_a}{x_a y_1 - x_1 y_a} u_b = 1. \quad (6)$$

It is non- S -trivial, so the same unit-equation theorem bounds the number of possible u_b ; explicitly $m \leq (ps)^{2D} + p^{4s-4}$. Multiplying by the cycle length yields the positive-characteristic branch of (1).

For a number field the proof is separate. A smallest good residue characteristic q_0 satisfies

$$q_0 < 12s \log(5s), \quad |\mathbf{k}(\mathfrak{p}_0)| + 1 \leq [12s \log(5s)]^D. \quad (7)$$

In the indifferent case, combine distinct tail reductions with the Morton–Silverman cycle bound (Theorem 6); this is absorbed by the second term of $\eta(0, D, s)$. In the attracting case the cycle has at most the residue-field size in (7). The same coordinate calculation (6), now with $a = 2$, lies in a two-variable group of rank $2s - 2$. The number-field unit theorem gives at most 2^{16s-8} values of u_b , hence $m \leq 2^{16s-8} + 3$. Together with (7), this is the first term of $\eta(0, D, s)$.

Toy case, hardest point, and scope

For $K = \mathbf{Q}$ and good reduction at every finite prime, $R_S^* = \{\pm 1\}$. The distance relations then allow only periods 1, 2, 3; indifferent tails have at most two points, while attracting tails have at most four points after passing to the cycle iterate. Corollary 1.2 concludes $|\mathcal{O}_\phi(P)| \leq 12$. This is the finite-unit shadow of the general argument.

The hard point is positive characteristic: Frobenius families such as $x + y = 1$ need not be finite. The proof must isolate a genuinely nontrivial equation before applying its quantitative theorem, including the purely inseparable case. Theorem 1 bounds one finite orbit, not the total number of preperiodic points. Corollary 1.1 derives a total-set bound only after fixing $\deg \phi$; the displayed constants are explicit but intentionally coarse.