

Vector bundles over a real elliptic curve

Indranil Biswas, Florent Schaffhauser

arXiv:1410.6845

<http://arxiv.org/abs/1410.6845v2>

The real form of the elliptic moduli space

A real elliptic curve is a complex elliptic curve X with an antiholomorphic involution σ ; its chosen origin need not lie in $X^\sigma = \text{Fix}(\sigma)$, which may be empty. Let $\mathcal{M}_X(r, d)$ be the coarse moduli variety of S-equivalence classes of semistable bundles, with real structure $E \mapsto \overline{\sigma^* E}$, and put

$$h = \gcd(r, d), \quad r' = r/h, \quad d' = d/h.$$

Main theorem (Biswas–Schaffhauser, Theorems 1.1 and 1.3). *As real algebraic varieties,*

$$\mathcal{M}_X(r, d) \simeq_{\mathbf{R}} \begin{cases} \text{Sym}^h(X), & X^\sigma \neq \emptyset, \\ \text{Sym}^h(X), & X^\sigma = \emptyset \text{ and } d' \text{ is odd}, \\ \text{Sym}^h(\text{Pic}_X^0), & X^\sigma = \emptyset \text{ and } d' \text{ is even}. \end{cases} \quad (1)$$

If $\mathcal{I}_X(r, d)$ parametrizes indecomposable bundles, then

$$\mathcal{I}_X(r, d) \simeq_{\mathbf{R}} \mathcal{M}_X(r', d') \simeq_{\mathbf{R}} \begin{cases} X, & X^\sigma \neq \emptyset, \\ X, & X^\sigma = \emptyset \text{ and } d' \text{ is odd}, \\ \text{Pic}_X^0, & X^\sigma = \emptyset \text{ and } d' \text{ is even}. \end{cases} \quad (2)$$

Although $X \simeq_{\mathbf{C}} \text{Pic}_X^0$, the two real forms need not be isomorphic when X^σ is empty; the parity switch in (1)–(2) is therefore genuine.

Ingredient I: Atiyah–Tu plus determinant

Every polystable representative of rank r , degree d is a direct sum $E_1 \oplus \cdots \oplus E_h$ of stable bundles of coprime rank r' and degree d' . Atiyah–Tu identify

$$\mathcal{M}_X(r, d) \longrightarrow \text{Sym}^h(\text{Pic}_X^{d'}), \quad [E_1 \oplus \cdots \oplus E_h] \longmapsto [\det E_1, \dots, \det E_h] \quad (3)$$

as a complex isomorphism. It is real because determinant commutes with $E \mapsto \overline{\sigma^* E}$.

The real Picard varieties provide the missing descent datum. If $X^\sigma \neq \emptyset$, a real basepoint identifies every Pic_X^m with X over $\mathbf{R}$. If $X^\sigma = \emptyset$, no real divisor of odd degree exists, and

$$\text{Pic}_X^m \simeq_{\mathbf{R}} \begin{cases} X, & m \text{ odd}, \\ \text{Pic}_X^0, & m \text{ even}. \end{cases} \quad (4)$$

Applying (4) to $m = d'$ in (3) proves (1). The hard conceptual point is exactly that one cannot silently choose a real origin when X^σ is empty.

Ingredient II: fixed points in the coprime case

When $h = 1$, every semistable bundle is stable and determinant identifies $\mathcal{M}_X(r, d)$ with Pic_X^d . A stable bundle fixed by the real involution admits, uniquely up to isomorphism, either a real structure $\tau^2 = 1$ or a quaternionic structure $\tau^2 = -1$, but not both. Combining this fact with the topological classification of such bundles gives Theorem 1.2:

- if $X^\sigma \neq \emptyset$, the fixed locus has one or two circle components, all parametrizing real bundles; the components are distinguished by $w_1(E^\tau)$;
- if $X^\sigma = \emptyset$ and d is odd, the fixed locus is empty;
- if $X^\sigma = \emptyset$ and d is even, it is two circles, one real and one quaternionic, interchanged by tensoring with a degree-zero quaternionic line bundle.

The determinant retains the real/quaternionic label in the last case because $\gcd(r, d) = 1$ forces r to be odd.

Ingredient III: make the indecomposable family real

Let F_h be Atiyah's unique indecomposable rank- h , degree-zero bundle with a nonzero section. It is built from the nonsplit extensions

$$0 \longrightarrow \mathcal{O}_X \longrightarrow F_h \longrightarrow F_{h-1} \longrightarrow 0. \quad (5)$$

Inductively, F_{h-1} is real and $H^1(X, F_{h-1}^*)$ is one-dimensional over $\mathbf{C}$; its real fixed line contains nonzero extension classes, all giving F_h with its real structure uniquely up to isomorphism. Thus F_h is canonically real even when X has no real points.

Atiyah–Tu's bijection

$$\mathcal{M}_X(r', d') \longrightarrow \mathcal{I}_X(r, d), \quad E' \longmapsto E' \otimes F_h \quad (6)$$

is consequently real, proving (2). It also shows that a fixed indecomposable bundle is real or quaternionic and transfers the preceding circle/emptiness classification with d replaced by d' . This is the proof mechanism behind Theorem 1.4.

Toy model and limitations

For $(r, d) = (2, 0)$, one has $\mathcal{M}_X(2, 0) \simeq_{\mathbf{R}} \text{Sym}^2(\text{Pic}_X^0)$, while every indecomposable is $L \otimes F_2$ and $\mathcal{I}_X(2, 0) \simeq_{\mathbf{R}} \text{Pic}_X^0$. Inside the symmetric square this is the diagonal $[L, L]$: it displays both the dimension drop from all S-equivalence classes to indecomposables and the role of (5).

The modular interpretation of every real point as a uniquely real or quaternionic bundle is asserted for the stable, hence coprime, case. For $h > 1$, the coarse symmetric product can also have fixed points made from conjugate pairs, so (1) alone is not a classification of individual real bundle structures. The paper treats vector bundles on genus-one curves; the unusually small dimension $\dim \mathcal{M}_X(r, d) = h$ and the Atiyah bundles are special to that setting.