The Dynamical Mordell–Lang Conjecture for Relatively Étale Systems over Products of Curves

Bhawesh Mishra

arXiv:2609.15121v1 (2026), 38 pages
http://arxiv.org/abs/2609.15121v1

Problem and setup

For an endomorphism $F\colon X \to X$, dynamical Mordell–Lang predicts that

$$S_F(x, V) := \{n \in \mathbb{N}_0 : F^n(x) \in V\}$$

is a finite union of arithmetic progressions. The paper studies a semiconjugacy $p\colon X \to B$ with $p \circ F = \beta \circ p$. Its *relative graph* is

$$\delta_F\colon X \longrightarrow B \times_{\beta,B,p} X, \qquad x \longmapsto (p(x), F(x)).$$

Calling F relatively ètale over β means that δ_F is ètale: the total map may inherit all ramification of the base, but contributes none in the relative directions.

Main theorem (Theorem 1.1)**.** *Let $B = \prod_{i=1}^{s} C_i$ be a finite product of smooth projective connected curves over $\mathbb{C}$ (the empty product is a point), and let $\beta\colon B \to B$ be any endomorphism. Let X be quasi-projective, $p\colon X \to B$ smooth, and $F\colon X \to X$ satisfy $p \circ F = \beta \circ p$. If F is relatively ètale over β, then $S_F(x, V)$ is a finite union of arithmetic progressions for every $x \in X(\mathbb{C})$ and every closed subvariety $V \subseteq X$.*

Taking $X = B$, $p = \mathrm{id}$, and $F = \beta$ makes the relative graph an isomorphism. Thus the theorem includes arbitrary endomorphisms of products of curves, with no restriction on the dimension or position of V and no genericity or postcritical hypothesis. This removes the possible infinite sparse remainder left by density-zero and gap-principle results.

Four proof ingredients

1. Simultaneous critical avoidance. The arithmetic core first treats a coordinatewise map $\Phi = (\phi_1, \ldots, \phi_g)$ on $(\mathbb{P}^1)^g$ over a number field. Theorem 2.3 finds a positive-density set of primes at which every chosen nonperiodic critical point remains nonperiodic. Locally, backward branches above a periodic critical reduction acquire inertia orbits of bounded number but increasingly divisible size. A Haar-measure estimate in a Frobenius coset then produces a simultaneous derangement in the joint preimage field; Chebotarev supplies the primes. Theorem 6.1 upgrades this to uniform avoidance: after one time M, every wandering coordinate orbit avoids the reductions of all critical points. An attracting periodic residue class necessarily contains a critical reduction, so a p-adic split-map theorem applies and writes the return set as finitely many suborbits.

2. From the base to the total space. Lemma 7.2 shows that relative ètaleness survives iteration and equivariant base change, and that ètaleness of the reduced base map along an orbit forces ètaleness of F there. Lemma 7.3 then interpolates an orbit of a smooth $\mathbb{Z}_\ell$-scheme whenever the reduced orbit is ètale: after passing to finitely many residue-class tails, a suitable iterate is congruent to the identity, Poonen interpolation produces analytic orbit maps, and Strassmann's theorem makes each zero set finite or an entire progression. Preperiodic base coordinates are removed separately by slicing into periodic residue classes.

3. Returning from algebraic to complex coefficients. For a complex system, Lemma 8.1 specializes all wandering base coordinates to algebraic wandering points only to select a good arithmetic reduction. Lemma 8.2 then embeds the *original* finitely generated coefficient field injectively into $\mathbb{Q}_\ell$ while prescribing that reduction. The injectivity, rather than the specialization, ensures that membership in V and hence the return set are unchanged.

4. Stable products of curves. Theorem 2.5 passes to an iterate and a stable image B_0, then constructs an equivariant surjection

$$(\mathbb{P}^1)^r \times Y \longrightarrow B_0, \qquad (\phi_1, \ldots, \phi_r, \eta),$$

where $\deg \phi_i \geq 2$ and η is surjective ètale. A dependency graph shows that transient curve coordinates disappear under iteration, while the positive-genus stable part has an ètale cover. Lemma 10.1 absorbs Y into the relative directions; pullback proves the result on the cover, and surjectivity plus passage from the iterate and orbit tail gives Theorem 1.1.

Roadmap, toy case, and limitation

For $X = B = (\mathbb{P}^1)^g$ and $p = \mathrm{id}$, the proof reduces visibly to critical avoidance followed by p-adic interpolation; the relative machinery is exactly what permits extra smooth fibers without adding ramification. The hardest step is the simultaneous Hasse principle: one Frobenius element must avoid fixed points in several backward trees at once. The method genuinely uses criticality—nontrivial local degrees create the growing inertia orbits— and the paper does not obtain the analogous simultaneous statement for arbitrary marked points.

Codex mathematical paper digest, 2026-10-26. Source paper should be consulted for full statements and proofs.