On vector bundles over moduli spaces trivial on Hecke curves

Indranil Biswas, Tomás L. Gómez

arXiv:2004.03223v2

http://arxiv.org/abs/2004.03223v2

Hecke curves as test curves

Let X be a smooth complex projective curve of genus $g \geq 3$, let $r \geq 2$, and let ξ be a line bundle whose degree d is coprime to r. Put

$$M = M_X(r, \xi),$$

the smooth projective moduli space of stable rank-r bundles with determinant ξ. It has $\mathrm{Pic}(M) \simeq \mathbb{Z}$ and is simply connected.

Tensoring every bundle by a fixed line bundle L gives an isomorphism

$$M_X(r, \xi) \xrightarrow{\sim} M_X(r, \xi \otimes L^r).$$

It preserves stability and carries Hecke curves to Hecke curves: tensoring an elementary-transform sequence by L gives the corresponding transform for $F \otimes L$. Since $\gcd(d, r) = 1$, the proof may therefore replace d by its unique representative with $0 < d < r$. This normalization is used in the Chern-class calculation below, while the theorem itself allows arbitrary degree coprime to r.

If F is a $(0,1)$-stable rank-r bundle—meaning that every proper subbundle G satisfies $\deg(G)/\operatorname{rk}(G) < (\deg(F)-1)/r$—with $\det F = \xi(x)$, the elementary transforms

$$0 \longrightarrow F_q \longrightarrow F \longrightarrow k_x \longrightarrow 0$$

are parametrized by $\mathbb{P}(F_x^\vee)$. Restricting this Hecke space to a line $\mathbb{P}^1 \subset \mathbb{P}(F_x^\vee)$ gives a *Hecke curve* in M, of degree r with respect to the ample generator.

Main theorem (Theorem 6.4)**.** *If $\mathcal{E}$ is a vector bundle on $M_X(r, \xi)$ whose restriction to every Hecke curve is trivial, then $\mathcal{E}$ is trivial.*

This is the moduli-space analogue of the line test on projective space. The analogy is useful but incomplete: Hecke curves through one fixed point do not sweep M, so the projective-space descent proof cannot simply be copied.

Ingredient 1: $c_1 = 0$ and semistability

Restriction along a Hecke curve injects $H^2(M, \mathbb{Z}) \simeq \mathbb{Z}$ into $H^2(\mathbb{P}^1, \mathbb{Z})$. Therefore $c_1(\mathcal{E}) = 0$. If $\mathcal{E}$ were unstable, take a positive-slope saturated subsheaf $\mathcal{V} \subset \mathcal{E}$. Its non-locally-free locus has codimension at least two, and Lemma 3.3 lets a general Hecke curve P avoid that locus. Since $\mathrm{Pic}(M) = \mathbb{Z}$, restriction preserves the sign of degree, so $\deg(\mathcal{V}|_P) > 0$. But the trivial bundle $\mathcal{E}|_P$ has no positive-degree subsheaf. This contradiction proves semistability.

Ingredient 2: coordinates for the second Chern class

Choose a Poincaré bundle $\mathcal{P}$ on $X \times M$, and let $p \in X$. Section 2 gives

$$H^4(M, \mathbb{Q}) = \left\langle f_2^2, b_2, f_3 \right\rangle_{\mathbb{Q}}, \qquad f_2 = c_2(\mathcal{P})/[X], \quad b_2 = c_2(\operatorname{End}\mathcal{P})/[p], \quad f_3 = c_3(\mathcal{P})/[X],$$

with $f_3 = 0$ when $r = 2$. Write $c_2(\mathcal{E}) = \alpha f_2^2 + \beta b_2 + \gamma f_3$. The proof constructs two complementary families of Hecke curves to kill the three coefficients.

Ingredient 3: a two-point Hecke family

Starting from a $(0, 2)$-stable bundle and modifying it at two fixed points produces

$$\psi : \mathbb{P}^{r-1} \times \mathbb{P}^{r-1} \longrightarrow M.$$

On a line in either factor with the other coordinate fixed, ψ is a Hecke curve. Applying the projective-space line test successively to the two factors shows that $\psi^*\mathcal{E}$ is trivial, hence $\psi^* c_2(\mathcal{E}) = 0$.

Normalize $\mathcal{P}$ so that $d' = \deg(\mathcal{P}|_{\{p\}\times M})$ satisfies $0 < d' < r$; equation (2.3) gives $dd' \equiv 1 \pmod{r}$. Let H_1, H_2 be the hyperplane classes of the two projective-space factors. The explicit pullbacks in (4.6) have $\psi^* b_2 = 0$, while the images of f_2^2 and f_3 separate the independent classes $(H_1 + H_2)^2$ and $H_1^2 + H_2^2$ when $r > 2$. Because the preceding tensor twist arranged $d, d' > 0$, the scalar $1 + dd'(r - 1)$ is positive. Thus $\alpha = \gamma = 0$; for $r = 2$, where $f_3 = 0$, the same computation gives $\alpha = 0$. Only βb_2 remains.

Ingredient 4: moving the modification point

Section 5 builds a second classifying map $\varphi : Z \to M$. Here Z fibres over a connected finite cover $T \to X$; the cover supplies an r-th root needed to keep the determinant fixed while the Hecke point moves. Every fibre of $Z \to T$ maps to a Hecke curve. Hence $\varphi^*\mathcal{E}$ descends to a bundle on the curve T, so its second Chern class vanishes. Before pullback, the calculation uses the positive generator $[\varpi] \in H^4(\mathbb{P}(Q^\vee), \mathbb{Z})$ and gives $c_2(\operatorname{End} F_p) = 2r[\varpi]$. The cover $q : Z \to \mathbb{P}(Q^\vee)$ has degree r^{2g}, so (5.8) gives, with $[\varpi_Z]$ the positive generator on Z,

$$\varphi^* b_2 = 2r^{\,2g+1}[\varpi_Z] \neq 0.$$

Pulling back $c_2(\mathcal{E}) = \beta b_2$ therefore forces $\beta = 0$.

Finish and scope

The semistable bundle $\mathcal{E}$ now has $c_1 = c_2 = 0$. Simpson's flatness result, in the form used in the paper, supplies a filtration with stable zero-Chern-class quotients and a flat holomorphic connection. Its monodromy representation is trivial because M is simply connected; therefore $\mathcal{E}$ is trivial.

The hardest step is not the final flatness argument but the detection of every direction in $H^4(M, \mathbb{Q})$: the fixed two-point family kills two coordinates, while the moving-point family detects the otherwise invisible b_2. The proof uses $g \geq 3$, coprimality of r and $\deg \xi$, and triviality on every Hecke curve. It does not assert that curves through one chosen point suffice.