Convex hulls of polynomial Julia sets

Malgorzata Stawiska

arXiv:2004.12521v1

http://arxiv.org/abs/2004.12521v1

A convex trapping region from the Julia set

Let $p \in \mathbb{C}[z]$ have degree $d \geq 2$. Write J_p for its Julia set, $K_p = \{z : (p^{\circ n}(z))_{n\geq 0} \text{ is bounded}\}$ for the filled Julia set, and

$$H_p = \operatorname{conv}(J_p) \subset \mathbb{C} \simeq \mathbb{R}^2.$$

Complete invariance gives $p^{-1}(J_p) = J_p$ and $p^{-1}(K_p) = K_p$, but convexification need not commute with a nonlinear map. The paper proves that it nevertheless preserves the direction needed for backward trapping.

Main theorem (Theorems 3.1, 3.6, and 3.7)**.** *For every complex polynomial p of degree $d \geq 2$,*

$$p^{-1}(H_p) \subseteq H_p.$$

If equality holds, then p is affinely conjugate to T_d, to $-T_d$, or to cz^d with $|c| = 1$. The standard representatives in this list do realize equality: their Julia sets are respectively the segment $[-1, 1]$ or the unit circle.

Thus a set defined by an ordinary Euclidean operation remembers strong dynamical information. The inclusion is universal; equality is rigid.

Ingredient 1: critical points cannot escape the hull

The decisive preliminary fact is Lemma 3.2: every zero of p' lies in H_p. If a critical point x_0 were outside H_p, separation supplies a closed half-plane $E \ni x_0$ disjoint from J_p. A refinement of the Gauss–Lucas theorem used in the paper says that any closed half-plane meeting the convex hull of the critical points is mapped surjectively onto $\mathbb{C}$. Hence $p(E) = \mathbb{C}$. For $w \in J_p$, this produces a preimage of w in E, contradicting $p^{-1}(w) \subset J_p$ and $E \cap J_p = \varnothing$.

For $p(z) = z^2 + c$, this step has a transparent toy model: J_p is centrally symmetric, so its unique critical point 0 is the midpoint of two Julia points. The half-plane argument is what survives in arbitrary degree.

Ingredient 2: all-fibres-in-B is a convex condition

For a closed convex $B \subset \mathbb{C}$ containing the zeros of p', set

$$C_B = \{w \in \mathbb{C} : p^{-1}(w) \subseteq B\}.$$

Lemma 2.10 proves that C_B is convex. Given $w_1, w_2 \in C_B$ and positive integers n_1, n_2, apply Gauss–Lucas to

$$P(z) = (p(z) - w_1)^{n_1}(p(z) - w_2)^{n_2}.$$

All zeros of P lie in B, so all zeros of P' do too. The remaining factor of P', after the familiar fibre and critical factors, is $p(z) - \frac{n_2 w_1 + n_1 w_2}{n_1 + n_2}$. Hence this rational convex combination belongs to C_B; continuity of roots makes C_B closed, so density gives every real convex combination.

Take $B = H_p$. Ingredient 1 makes the lemma applicable. Complete invariance gives $J_p \subset C_{H_p}$, and convexity then gives $H_p \subset C_{H_p}$, which is exactly $p^{-1}(H_p) \subset H_p$.

Equality: segment or disk

Always $K_p \subseteq H_p$: in one complex variable, K_p is the holomorphic hull of J_p, and the holomorphic hull lies in the ordinary convex hull. Conversely, if $p^{-1}(H_p) = H_p$, every forward orbit starting in H_p stays in the compact set H_p, so $H_p \subseteq K_p$. Thus equality forces $H_p = K_p$.

If this convex set has empty interior, then $J_p = K_p = H_p$ is a compact segment. An affine change sends it to $[-1, 1]$, and complete invariance of that segment characterizes T_d and $-T_d$. If the interior is nonempty, $J_p = \partial H_p$ is a rectifiable Jordan curve, hence has Hausdorff dimension 1. Hamilton's theorem says that a rational Julia Jordan curve has dimension > 1 unless it is a circle or line; the polynomial case here leaves a circle. After normalizing it to the unit circle, complete invariance characterizes cz^d, $|c| = 1$.

The hardest point is the convexity of C_B: it converts information about critical points into simultaneous control of every root in every fibre. The classification additionally depends on external rigidity results for invariant segments/circles and Julia Jordan curves; the note does not replace those inputs.