A Torelli theorem for moduli spaces of parabolic vector bundles over an elliptic curve

Thiago Fassarella, Luana Justo

arXiv:2008.05229v2 (2020)
http://arxiv.org/abs/2008.05229v2

Can a parabolic moduli space remember its curve?

Let C be a complex elliptic curve, let $S = p_1 + \cdots + p_n$ be a reduced divisor with $n \geq 3$, and fix $w \in C$. A rank-two quasiparabolic bundle consists of a bundle E together with a line $V_i \subset E_{p_i}$ at every marked point. For the central weight $a_F = (1/2, \ldots, 1/2)$, write $\mathcal{M}_{a_F}(C, S, w)$ for the moduli space of μ_{a_F}-semistable such bundles with determinant $\mathcal{O}_C(w)$.

Main theorem (Theorems 1.1 and 4.11)**.** *Let C, C' be elliptic curves, and let $S \subset C$, $S' \subset C'$ be reduced divisors of the same degree $n \geq 3$. If*

$$\mathcal{M}_{a_F}(C, S, w) \simeq \mathcal{M}_{a_F}(C', S', w'),$$

then there is an isomorphism $C \simeq C'$ carrying S to S'.

Thus the abstract projective moduli space determines the marked curve, even though neither C nor the parabolic points are visibly part of a point of that space. The condition $n \geq 3$ is sharp for this argument and for the statement as given: when $n = 2$, the moduli space is always $\mathbb{P}^1 \times \mathbb{P}^1$.

Spectral setup and its discriminant

Let $\mathcal{H}_{a_F}$ be the moduli space of trace-free semistable parabolic Higgs bundles $(E, \mathbf{V}, \theta)$ with the same determinant. Its stable locus contains $T^*\mathcal{M}^s_{a_F}$ as an open subset. The full Hitchin map and its cotangent restriction are

$$h : \mathcal{H}_{a_F} \longrightarrow \Sigma := H^0\big(C, \omega_C^{\otimes 2}(S)\big), \quad (E, \mathbf{V}, \theta) \longmapsto \det\theta, \qquad h^\circ := h|_{T^*\mathcal{M}^s_{a_F}}.$$

Put $P = \mathrm{Tot}(\omega_C(S))$, let $q : P \to C$ be its projection, and let s be the tautological section of $q^*\omega_C(S)$. For $a \in \Sigma$, the spectral double cover $C_a = \{s^2 + q^*a = 0\} \subset P$ has projection $q_a = q|_{C_a}$. Its singular locus in the Hitchin base is

$$\mathcal{D} = \widetilde{\mathcal{D}} \cup H_1 \cup \cdots \cup H_n, \quad H_i = H^0\big(\omega_C^{\otimes 2}(S - p_i)\big),$$

where $\widetilde{\mathcal{D}} = \bigcup_{p \in C} H^0(\omega_C^{\otimes 2}(S - 2p))$ records sections with a double zero.

The local toy model is $y^2 + a(x) = 0$: two sheets meet when a develops a double zero. At a parabolic point the allowed pole and the nilpotent residue shift this condition into the distinguished hyperplane H_i. The full proof keeps track of both types of singularity, because the nonlinear component $\widetilde{\mathcal{D}}$ remembers the curve while the hyperplanes remember its marked points.

Ingredient 1: the discriminant already contains (C, S)

Since $n \geq 3$, $\omega_C^{\otimes 2}(S)$ is very ample and embeds

$$C \hookrightarrow \mathbb{P}(\Sigma^\vee), \qquad p \longmapsto H^0(\omega_C^{\otimes 2}(S - p)).$$

Proposition 3.8 identifies the projective dual of $\mathbb{P}\widetilde{\mathcal{D}}$ with this embedded C, while the dual of each $\mathbb{P}H_i$ is precisely p_i. Hence knowing the pair $(\Sigma, \mathcal{D})$ reconstructs (C, S). The remaining problem is to recover this pair from the moduli space alone.

Ingredient 2: recover the Hitchin base intrinsically

The subtle step is Proposition 4.7. Inside the stable parabolic-Higgs moduli space, the complement of $T^*\mathcal{M}^s_{a_F}$ has codimension at least $n/2$, hence at least two. To prove this, Proposition 4.2 classifies, up to elementary transformations, the indecomposable parabolic bundles that fail central-weight stability. Decomposable bundles are reduced to $E = L \oplus L^{-1}(w)$ with $\deg L = 1$; one exceptional indecomposable case cannot support a semistable Higgs field because every such field leaves the destabilizing line invariant. Dimension counts for the remaining decomposable and indecomposable families give the codimension bound.

Hartogs extension now forces every regular function on $T^*\mathcal{M}^s_{a_F}$ to extend across that complement. For a general Hitchin parameter, the spectral correspondence identifies the fiber with a Prym variety, so all of its regular functions are constant. Proposition 4.9 therefore identifies the affinization

$$\operatorname{Spec} \mathbb{C}[T^*\mathcal{M}^s_{a_F}] \simeq \Sigma,$$

and its canonical map with h°, up to an automorphism of Σ. This is why the Hitchin base is intrinsic to the original moduli space rather than an extra structure chosen in the proof.

Ingredient 3: detect singular spectral curves by fiber geometry

Proposition 3.10 separates the two generic fibers of the full map h. If $a \notin \mathcal{D}$, then C_a is smooth and integral of genus $n+1$, and $h^{-1}(a)$ is the fixed-determinant Prym

$$\{M \in \operatorname{Pic}^{n+1}(C_a) : \det(q_{a*}M) = \mathcal{O}_C(w)\},$$

an abelian variety. For general $a \in \mathcal{D}$, the spectral curve has one node and its compactified Jacobian, hence the full fixed-determinant fiber, is uniruled. The fibers of h°, which are the fibers available intrinsically from the cotangent bundle, are the corresponding open subsets. Thus outside $\mathcal{D}$ they lie in abelian varieties and contain no rational curves, whereas over a general point of $\mathcal{D}$ they are open subsets of uniruled varieties. This is the fiber-geometric distinction used in the proof of Theorem 4.11 to recognize $\mathcal{D}$.

Proof roadmap and scope

An isomorphism of the two parabolic-bundle moduli spaces restricts to their stable loci and induces an isomorphism of cotangent bundles. Affinization then gives an isomorphism $\xi : \Sigma \to \Sigma'$. Fiberwise scalar dilation makes ξ $\mathbb{C}^*$-equivariant, hence linear. The abelian-versus-uniruled dichotomy forces $\xi(\mathcal{D}) = \mathcal{D}'$; projective duality from Ingredient 1 finally sends (C, S) to (C', S').

The hardest technical work is the codimension estimate that legitimizes affinization and Hartogs extension; it depends on the explicit classification of unstable underlying parabolic bundles and on the central weight. The theorem is over $\mathbb{C}$, in rank two, with degree-one fixed determinant and weights a_F. Its conclusion recovers the marked pair (C, S), not the chosen determinant point w; indeed twisting by a square root identifies the moduli spaces for different degree-one determinants on a fixed elliptic curve.