p-Adic interpolation of orbits under rational maps

Jason P. Bell, Xiao Zhong

arXiv:2202.01673v1 (3 February 2022)

http://arxiv.org/abs/2202.01673v1

The interpolation problem

Let L be a field of characteristic zero, let $h : \mathbb{P}^1 \to \mathbb{P}^1$ be a rational self-map defined over L, and fix $c \in \mathbb{P}^1(L)$. A single power series in the iteration number cannot usually parametrize the whole orbit: a preperiodic orbit, for example, may cycle through several values, whereas a convergent p-adic series constant on infinitely many integers is constant. The right object is therefore a tail of the orbit, separated into congruence classes.

Main theorem (Bell–Zhong, Theorem 1.1)**.** *There is a finitely generated extension $K/\mathbb{Q}$, over which h and c are defined, and an infinite set of inequivalent non-archimedean completions $K_{\mathfrak{p}}$ with the following property. For every such completion there is an integer $a = a(\mathfrak{p}) > 0$, and for each $i \in \{0, \dots, a-1\}$ a series $h_i(t) \in K_{\mathfrak{p}}[[t]]$, convergent on the closed unit disc, such that*

$$h^{an+i}(c) = h_i(n) \qquad \text{for every sufficiently large integer } n.$$

Neither the initial tail nor the dependence of a on $\mathfrak{p}$ is suppressed. What is striking is the abundance of favorable completions for an arbitrary one-dimensional rational orbit; at a fixed prime the orbit could instead lie in an attracting or superattracting regime that has no analytic parametrization by the iteration number.

How favorable completions are manufactured

The degree-one and eventually periodic cases are immediate from known étale interpolation and constancy on progressions. In the remaining case, pass to an iterate of h having at least four non-superattracting fixed points and conjugate one of them to infinity. Thus $h = p/q$, with $\gcd(p, q) = 1$ and $\deg p = \deg q + 1$. Three finite fixed points $\gamma_1, \gamma_2, \gamma_3$ may be chosen away from the critical points.

The authors build a finitely generated ring R containing the coefficients, the relevant roots, and the necessary inverses. They write $h^n(c) = A_n/B_n$ using explicit homogeneous recurrences. Lemma 2.1 says $A_nR + B_nR = R$, so reduction never makes both coordinates vanish.

The central arithmetic step is Lemma 2.2. Unless the orbit is eventually periodic, there is a Zariski-dense set of maximal ideals $\mathfrak{p} \subset R$ such that, for some n, the reduction of A_n/B_n is a fixed point of h, its denominators are nonzero, and its multiplier is nonzero. Here is the rigidity behind this assertion. If no such primes remained after a localization, then all three quantities $A_n - \gamma_j B_n$ would be units. They satisfy a fixed three-term linear relation. The S-unit theorem gives only finitely many such triples up to scale, so two orbit values must coincide—contradicting nonperiodicity. This argument is the technical bottleneck: it simultaneously keeps the reduced orbit defined, forces recurrence, and avoids critical reduction.

From a congruence to an analytic orbit

Restrict the preceding maximal ideals to the regular locus. Their residue fields are finite, and maximal-ideal order supplies a discrete valuation on the corresponding completion. Finiteness lets one choose a tail point c_*, an iterate $H = h^a$, and the residue-characteristic prime $\pi \geq 5$, with $|\pi| < 1$, so that

$$H(c_*) \equiv c_* \pmod{\pi^2}, \qquad H'(c_*) \equiv 1 \pmod{\pi}.$$

Lemma 2.3 rescales the local dynamics to

$$F(x) = \pi^{-1}\big(H(c_* + \pi x) - c_*\big), \qquad F(x) \equiv x \pmod{\pi},$$

with F in the Tate algebra of the closed unit disc. Poonen's theorem then produces $G(x,t)$ in a two-variable Tate algebra with $F^n(x) = G(x,n)$. Evaluating at the rescaled starting point interpolates the H-orbit; undoing the tail and the iterate gives the a series in the theorem.

A transparent local toy model is $H(x) = x + p^2$ over $\mathbb{Q}_p$, with $p \geq 5$ and $c_* = 0$. Then $F(x) = p^{-1}H(px) = x + p$, so $H^n(0) = p^2 n$ is already analytic in n. This displays the final congruence-to-identity mechanism, but omits the hard global task of finding infinitely many places where an arbitrary orbit admits such coordinates.

The Dynamical Mordell–Lang consequence

Let X be a complex quasiprojective variety, let $g : X \to X$ be étale, and set $\Phi = (h, g)$ on $\mathbb{P}^1 \times X$, with $h \in \mathbb{C}(x)$. Corollary 1.3 states that for every point z and closed subvariety $Y \subseteq \mathbb{P}^1 \times X$,

$$\{n \geq 0 : \Phi^n(z) \in Y\}$$

is a finite union of infinite arithmetic progressions together with a finite set. Enlarge the arithmetic model R to include X and g, and choose a favorable maximal ideal inside the required smooth model. The main theorem interpolates the $\mathbb{P}^1$-coordinate, while étaleness gives analytic coordinates for the X-orbit. On a common iterate, the equations of Y become finitely many one-variable p-adic analytic functions. By Strassmann's theorem, their common integer zero set is either finite or eventually all of that progression. Taking all residue classes gives the conclusion.

Scope and limitations

The same interpolation conclusion extends to self-maps of curves: after normalization, genus zero is the theorem, genus one is covered by étale methods, and in genus at least two every endomorphism has finite order. Higher dimension is different. The map $(x, y) \mapsto (x + 1, y(x + 1))$ has orbit $(n, n!)$ from $(0, 1)$, which cannot be p-adically interpolated at any prime because $|n!|_p \to 0$. Even simultaneous interpolation for two arbitrary rational maps on $(\mathbb{P}^1)^2$ remains open. Thus the split Dynamical Mordell–Lang corollary uses exactly the combination available here: one arbitrary rational coordinate and an étale complementary system.