Toric varieties with ample tangent bundle

Kuang-Yu Wu

arXiv:2210.01366v1 (4 October 2022)

https://arxiv.org/abs/2210.01366v1

The toric form of Mori's theorem

Work over an algebraically closed field k. Recall that a vector bundle E is ample when $\mathcal{O}_{\mathbb{P}(E)}(1)$ is ample. Wu proves the following toric classification without bend-and-break.

Main theorem (Wu, Theorem 1.1)**.** *Let X be an n-dimensional smooth projective toric variety over k. If its tangent bundle T_X is ample, then*

$$X \cong \mathbb{P}^n_k.$$

The point is not a new classification beyond Mori's theorem, but a direct dictionary between positivity and convex geometry. Choose any ample torus-invariant divisor D. The pair (X, D) determines a full dimensional lattice polytope P, whose normal fan is the fan of X. Smoothness makes P simple: exactly n edges leave every vertex. The proof shows that ampleness of T_X forces every two-dimensional face of P to be a triangle; simplicity then forces P itself to be an n-simplex.

From an invariant curve to integers

Let τ be a codimension-one cone of the fan, $C_\tau \cong \mathbb{P}^1$ the corresponding invariant curve, and σ, σ' the two maximal cones meeting along τ. Smoothness gives primitive generators

$$\sigma = \langle v_1, \ldots, v_{n-1}, v_n \rangle, \qquad \sigma' = \langle v_1, \ldots, v_{n-1}, v'_n \rangle$$

and dual bases $u_1, \ldots, u_n$ and $u'_1, \ldots, u'_n$ of the character lattice M. These dual bases are precisely the associated characters of T_X on the two affine toric charts.

Only u_i and u'_i have difference in the rank-one lattice $\tau^\perp \cap M$. Writing

$$u_i = u'_i + a_i u_n, \qquad a_i \in \mathbb{Z},$$

with $u'_n = -u_n$ and hence $a_n = 2$, the equivariant gluing calculation gives Proposition 3.3:

$$T_X|_{C_\tau} \cong \bigoplus_{i=1}^{n} \mathcal{O}_{\mathbb{P}^1}(a_i).$$

A toric vector bundle is ample exactly when all of its restrictions to invariant curves are ample. Thus ampleness of T_X says $a_i > 0$ for every wall τ and every i.

From splitting degrees to angles

Let $p_\sigma, p_{\sigma'}$ be the vertices of P joined by the edge corresponding to τ. For each $j < n$, that edge and the edge directions u_j, u_j' lie in a common two-face A_j. Proposition 4.1 translates the integer a_j into the two angles adjacent to the shared edge:

$$\begin{array}{l|l} a_j > 0 & \theta(p_\sigma, A_j) + \theta(p_{\sigma'}, A_j) < \pi \\ a_j = 0 & \theta(p_\sigma, A_j) + \theta(p_{\sigma'}, A_j) = \pi \\ a_j < 0 & \theta(p_\sigma, A_j) + \theta(p_{\sigma'}, A_j) > \pi. \end{array}$$

The proof is the elementary geometry of a trapezoid formed by the two parallel edge directions. Although the individual angles depend on a choice of Euclidean coordinates on $M_\mathbb{R}$, the trichotomy above does not. Therefore T_X ample implies that the sum of the two endpoint angles along every edge of every two-face is strictly below π.

This step is the technical hinge: the splitting calculation must match the characters across τ correctly before the sign of a_j can be read as a convex-geometric inequality. Once that bridge is in place, the remaining argument is short.

The polygon count and the simplex

Take a two-face A with cyclically ordered vertices $p_1, \dots, p_m$. Adding the strict inequality along all m edges gives

$$m\pi > \sum_{i=1}^{m} \big(\theta(p_i, A) + \theta(p_{i+1}, A)\big) = 2\sum_{i=1}^{m} \theta(p_i, A) = 2(m-2)\pi.$$

Hence $m < 4$, so every two-face is a triangle.

Now fix a vertex q_0 and its n neighbors $q_1, \dots, q_n$. The two edges q_0q_i, q_0q_j lie in a two-face, which is triangular, so q_i and q_j are adjacent. Each q_i already has its full complement of n neighbors among the other q_j's and q_0. Thus the connected edge graph has only these $n+1$ vertices: P is an n-simplex. Its smooth normal fan is the fan of $\mathbb{P}^n$, proving the theorem.

A surface test and the scope

The examples display all three signs. For $\mathbb{P}^2$, an invariant line has $T_X|_C \cong \mathcal{O}(1) \oplus \mathcal{O}(2)$; the adjacent-angle sum is $< \pi$. For $\mathbb{P}^1 \times \mathbb{P}^1$ the splitting $\mathcal{O}(0) \oplus \mathcal{O}(2)$ makes T_X nef but not ample and gives equality. On the Hirzebruch surface $\mathbb{F}_1$, the splitting $\mathcal{O}(-1) \oplus \mathcal{O}(2)$ gives an angle sum $> \pi$. The argument depends on the established invariant-curve positivity criterion for toric vector bundles and on projectivity, which supplies P; it is a toric reproof, not a replacement for Mori's general theorem.