Torelli theorem for moduli stacks of vector bundles and principal G-bundles

David Alfaya, Indranil Biswas, Tomás L. Gómez, Swarnava Mukhopadhyay

arXiv:2310.07666v2 (25 November 2024)
http://arxiv.org/abs/2310.07666v2

What the stack remembers

Let X be an irreducible smooth complex projective curve, let $g(X) \geq 2$, fix $r \geq 2$, and choose a line bundle ξ. Write $\mathcal{M}(X, r, \xi)$ for the full moduli stack whose objects are pairs

$$(E, \varphi), \qquad \operatorname{rk} E = r, \qquad \varphi\colon \det E \xrightarrow{\sim} \xi,$$

with arrows compatible with φ. It includes unstable bundles. This is crucial: the stabilizers and intrinsic substacks retain data that the coarse moduli scheme can lose.

Main theorem (Alfaya–Biswas–Gómez–Mukhopadhyay, Theorem 5.2)**.** *Let X, X' be irreducible smooth complex projective curves of genera at least 2, let $r, r' \geq 2$, and fix line bundles ξ, ξ'. If*

$$\mathcal{M}(X, r, \xi) \simeq \mathcal{M}(X', r', \xi'),$$

then $r = r'$ and $X \simeq X'$.

There is no coprimality condition on rank and determinant degree. The point is not simply to read off dimensions: the proof must reconstruct the curve even when distinct curves have isomorphic coarse spaces.

The Hitchin reconstruction

First remove three low-dimensional pairs $(g, r) = (2, 2), (2, 3), (3, 2)$. The simple objects are intrinsically the objects with minimal finite stabilizer, the scalar group μ_r. The degree-zero part of the simple locus's cotangent complex is the stack $\mathcal{N}^{\mathrm{simp}}$ of trace-free Higgs bundles. Its affinization recovers the Hitchin map

$$H\colon \mathcal{N}^{\mathrm{simp}} \longrightarrow W = \bigoplus_{k=2}^{r} H^0(X, K_X^{\otimes k}).$$

The natural $\mathbb{C}^*$-action has weights $2, \ldots, r$. Hence an isomorphism of stacks induces an equivariant isomorphism of Hitchin bases; the length of the weight filtration recovers r, and its highest-weight summand is $W_r = H^0(X, K_X^{\otimes r})$.

The discriminant $D \subset W$, where the spectral curve is singular, is also intrinsic. A map $\mathbb{P}^1 \to \mathcal{N}^{\mathrm{simp}}$ whose image contains two non-isomorphic objects is constant under H and must land over D: away from D, the Hitchin fiber is a Prym variety and contains no rational curve. Conversely, over a general point of D, the one-nodal spectral curve has a compactified Jacobian fibered by nodal rational curves. A codimension estimate keeps a complete such curve inside the cotangent locus, and an explicit family lifts it to the stack. Thus rational curves recover a dense open part of D, hence all of D.

Finally put

$$C = \mathbb{P}(D \cap W_r) \subset \mathbb{P}(W_r).$$

A projective-duality result used by the authors identifies $C^\vee \simeq X$. This completes the Torelli argument outside the three exceptional pairs. The hardest step is the converse rational-curve construction: the codimension bound must be strong enough that the one-dimensional curves in a general singular Hitchin fiber avoid the bad locus.

Recovering semistability from the stack

The second mechanism descends from the full stack to its usual projective moduli scheme without choosing semistability externally. A filtration is a map $\Theta = [\mathbb{A}^1/\mathbb{G}_m] \to \mathcal{M}$. The weight of the determinant-of-cohomology line bundle on its special fiber is the Hilbert–Mumford expression for slope semistability. Positive multiples therefore select precisely the slope-semistable bundles; the zero class selects everything and negative multiples select nothing. Consequently any line bundle whose semistable locus is nonempty and proper recovers the same intrinsic substack $\mathcal{M}^{\mathrm{ss-vb}}(X, r, \xi)$.

Its good moduli space is the familiar projective scheme $M^{\mathrm{ss-vb}}(X, r, \xi)$. A stack isomorphism therefore descends to an isomorphism of these schemes, allowing known scheme-level Torelli theorems to handle many cases.

Why the exceptional stack still works

The illuminating case is $g = 2, r = 2$ with even determinant degree. After twisting, one reduces to trivial determinant. The coarse space is $\mathbb{P}^3$ for every genus-two curve, so it cannot remember X. The stack does: its non-simple semistable objects project exactly onto the strictly semistable locus

$$K(X) = J(X)/\{\pm 1\}, \qquad [L] \longmapsto [L \oplus L^{-1}].$$

From the smooth part of this Kummer quotient one canonically recovers the double cover $J(X) \setminus J(X)[2] \to K(X)^{\mathrm{sm}}$, then its abelian compactification $J(X)$. The ample generator of the coarse moduli space recovers the canonical polarization, so classical Torelli recovers X.

The dimension formula $\dim M^{\mathrm{ss-vb}} = (r^2 - 1)(g - 1)$ isolates dimensions $3, 6, 8$. The dimension-three case uses the Kummer argument (or odd-degree Torelli), dimension six uses the known genus-three rank-two theorem, and dimension eight uses the Brauer group to distinguish determinant degree coprime to 3 from degree divisible by 3, followed by the appropriate known Torelli theorem. All larger cases use the Hitchin argument.

Principal bundles and scope

For an algebraic connected reductive complex group G, let $\mathcal{M}_G^d(X)$ be the component indexed by $d \in \pi_1(G)$. Lemma 3.7 intrinsically recovers Ramanathan semistability as the intersection of all nonempty semistable loci defined by rational line bundles; one such line bundle already realizes it. After passing to good moduli spaces, the cited principal-bundle Torelli theorem gives

$$\mathcal{M}_G^d(X) \simeq \mathcal{M}_{G'}^{d'}(X'), \quad g(X), g(X') \geq 3 \quad \Longrightarrow \quad X \simeq X'.$$

This corollary does not assert $G \simeq G'$. Likewise, the main theorem does not claim to recover the determinant line bundle itself. Everything is over $\mathbb{C}$, and several final steps deliberately invoke established Torelli results for coarse moduli spaces; the new work is the intrinsic passage from stack geometry to the structures those results need.