On the moduli space of simple sheaves on singular K3 surfaces

Barbara Fantechi, Rosa M. Miró-Roig

arXiv:2403.00735v2 (14 January 2025)

http://arxiv.org/abs/2403.00735v2

The singular K3 setting

Here a K3 surface is a projective, Gorenstein, reduced, connected complex surface X with $H^1(X, \mathcal{O}_X) = 0$ and $\omega_X \simeq \mathcal{O}_X$; it need not be normal or irreducible. A coherent sheaf F is *perfect* if it has a finite locally free resolution, and *simple* if $\mathrm{End}(F) = \mathbb{C}$. Write $\mathrm{PSpl}(r; L, c_2)$ for the algebraic space of perfect simple sheaves of rank $r \geq 1$, determinant L, and second Chern class c_2.

A pair (X, L) is smoothable when it occurs as the special fiber of a flat projective family over a smooth connected curve whose other fibers are smooth, and L extends to the total space. This condition on the *pair*, rather than only on X, is what transports the closedness of Mukai's form.

Main theorem (Fantechi–Miró-Roig, Theorem 1.1)**.** *The space* $\mathrm{PSpl}(r; L, c_2)$ *is smooth of dimension*

$$2rc_2 - (r-1)L^2 - 2r^2 + 2.$$

Serre duality supplies it with a natural nowhere-degenerate algebraic two-form α*. If* (X, L) *is smoothable, then* $d\alpha = 0$*, so* $\mathrm{PSpl}(r; L, c_2)$ *is a symplectic algebraic space.*

For smooth X, every coherent sheaf is perfect and the constant family is a smoothing; the theorem recovers the familiar Mukai picture. On a singular surface, perfectness is the exact hypothesis that keeps the required duality available.

Deformations, duality, and the two-form

At $[F]$, fixed-determinant deformation theory has tangent and obstruction spaces

$$\mathrm{Ext}^1(F, F)_0, \qquad \mathrm{Ext}^2(F, F)_0,$$

where the subscript denotes the kernel of trace. Perfect Serre duality on the Gorenstein surface identifies $\mathrm{Ext}^2(F, F)$ with $\mathrm{Hom}(F, F)^*$. Simplicity makes both one-dimensional; the trace splitting identifies their scalar summand with $H^2(\mathcal{O}_X) \simeq \mathbb{C}$. Hence the trace-free obstruction space vanishes, proving smoothness. Riemann–Roch gives the displayed dimension.

The form at F is the pairing

$$\mathrm{Ext}^1(F, F) \otimes \mathrm{Ext}^1(F, F) \xrightarrow{\smile} \mathrm{Ext}^2(F, F) \xrightarrow{\mathrm{tr}} H^2(\mathcal{O}_X) \simeq \mathbb{C}.$$

Graded commutativity makes it alternating and Serre duality makes it nondegenerate. Using a universal sheaf on the fixed-determinant Deligne–Mumford stack, relative Ext bundles, and étale descent, the pointwise pairings assemble into a form on the coarse algebraic space.

The technically new point is closedness. For a smoothing $\pi\colon \mathcal{X} \to B$, the relative moduli space is smooth over B, and the same cup–trace construction produces a relative form α_B. Over the dense open set of smooth K3 fibers, Mukai's theorem gives $d_\pi \alpha_B = 0$. This relative three-form is a regular section of a vector bundle, so vanishing on the dense open set forces vanishing everywhere. Restriction to the special fiber proves $d\alpha = 0$.

Lagrangians transported between moduli spaces

Two companion results show that familiar constructions survive the singular setting.

For a perfect simple vector bundle F with $H^0(F) = H^0(F^*) = 0$, choose $V^* \subset H^1(F^*)$, $\dim V = v$, and form

$$0 \longrightarrow V \otimes \mathcal{O}_X \longrightarrow E \longrightarrow F \longrightarrow 0.$$

On the indicated open locus this gives a locally closed embedding from a Grassmann bundle into $\mathrm{PSpl}(r+v; L, c_2)$. Theorem 1.2 says that, for every smooth irreducible locally closed subspace $\mathcal{L}$, its induced image is half-dimensional exactly when $\mathcal{L}$ is, and is isotropic (respectively Lagrangian) exactly when $\mathcal{L}$ is. No smoothing assumption is needed here.

For a globally generated F with $H^1(F) = 0$, choose $W \subset H^0(F)$, $w = \dim W \geq r + 2$, with surjective evaluation, and take the generalized syzygy bundle

$$0 \longrightarrow S \longrightarrow W \otimes \mathcal{O}_X \longrightarrow F \longrightarrow 0.$$

Now S has invariants $(w - r, -L, L^2 - c_2)$. Theorem 1.3 gives the same equivalence for half-dimensionality; when (X, L) is smoothable, it also gives the isotropic/Lagrangian equivalence. Its proof again spreads the relevant two-form identity over a smoothing and specializes from the dense smooth locus.

When the polarization smooths

The paper makes the main hypothesis testable. Suppose X has isolated local-complete-intersection singularities and L is ample. Let

$$0 \longrightarrow \Omega_X \longrightarrow \mathcal{P} \longrightarrow \mathcal{O}_X \longrightarrow 0$$

be the Atiyah extension of L, and put $\mathcal{T}^1_X = \mathcal{E}xt^1(\Omega_X, \mathcal{O}_X)$. Theorem 5.12 states that (X, L) is polarized smoothable if

$$H^2(X, T_X) = 0, \qquad \mathrm{Ext}^1(\mathcal{P}, \mathcal{O}_X) \twoheadrightarrow H^0(X, \mathcal{T}^1_X).$$

The first condition kills global obstructions; the second lets a global deformation of the pair realize simultaneous local smoothing directions. The obstruction group $\mathrm{Ext}^2(\mathcal{P}, \mathcal{O}_X)$ then vanishes, so the chosen first-order smoothing lifts formally, and algebraization produces a projective smoothing. Proposition 5.18 verifies these conditions for K3 surfaces with only ADE singularities.

The scope matters: the two-form is always nondegenerate, but its closedness is asserted only under smoothability; the moduli space need not be proper or nonempty. The Lagrangian constructions concern the specified open loci of vector bundles, not all perfect sheaves. These restrictions are precisely where singular geometry differs from the smooth K3 case.