Arakelov-Green's functions for dynamical systems on projective varieties

Nicole R. Looper

arXiv:2404.06981v1 (10 April 2024)

http://arxiv.org/abs/2404.06981v1

A global Green-function inequality in any dimension

Let K be a product formula field with normalized absolute values $|\cdot|_v$, and let (X, ϕ, L) be a polarized dynamical system: X/K is projective of positive dimension, L is ample, and $\phi^* L \simeq L^{\otimes d}$ for $d \geq 2$. After replacing L by a power, embed $X \hookrightarrow \mathbb{P}^N$ so that ϕ extends to a degree-d morphism $f\colon \mathbb{P}^N \to \mathbb{P}^N$.

Choose a homogeneous lift $F\colon \mathbb{A}^{N+1} \to \mathbb{A}^{N+1}$. At a place v, write $\mathcal{K}_v$ for its homogeneous filled Julia set, $X_v = X \times_K \mathbb{C}_v$, and $\pi_v\colon \mathbb{A}^{N+1}_{\mathbb{C}_v} \setminus \{0\} \to \mathbb{P}^N_{\mathbb{C}_v}$; put $E_v = \mathcal{K}_v \cap \pi_v^{-1}(X_v)$. If $B_n = (\eta_j)$ is a basis of $H^0(X, \iota^* \mathcal{O}(n))$, of dimension $c(n)$, set

$$d_{B_n}(E_v) = \max_{\widetilde{P}_1, \ldots, \widetilde{P}_{c(n)} \in E_v} \left| \det\big(\eta_j(\widetilde{P}_i)\big) \right|_v^{1/(nc(n))}.$$

This is a basis-dependent transfinite diameter. After the paper passes to an extension and normalizes the lift so that its resultant term vanishes, its negative logarithm is the extremal value of the generalized averaged Arakelov–Green function.

Main theorem (Looper, Theorem 3.3)**.** *For every choice of the bases B_n, there is $C = C(f)$ such that, for all $n \geq 2$,*

$$\sum_{v \in M_K} \log d_{B_n}(E_v) \leq \frac{C \log n}{n}.$$

Changing B_n multiplies every determinant by the same scalar over K; the product formula makes the sum independent of that change. This is why the assertion is non-vacuous globally even though a single local basis can be rescaled at will.

The special basis and the decisive estimate

The coordinate forms of the iterates $F^{(k)}$, raised to powers between 1 and $d-1$, generate a large collection $\mathcal{G}$. For $n \geq d(N+1)$, Macaulay elimination is applied recursively: every homogeneous form of degree n becomes a linear combination of products of only $O_{d,N}(\log n)$ members of $\mathcal{G}$, times a form of bounded degree. Restricting a maximal independent subcollection gives the special basis $H(n)$; smaller degrees use the standard monomial basis. No uniform rule identifies which redundant vectors to discard; the adelic basis-independence above is what makes this harmless.

On $\mathcal{K}_v$, iterated coordinate forms remain bounded. Thus each column of the $H(n)$-evaluation matrix has norm at most $\exp(C \log n)$. Hadamard's determinant inequality then gives

$$\log \left| \det\big(\eta_j(\widetilde{P}_i)\big) \right|_v \leq C \max\{\log R_v, 1\} (\log n) c(n),$$

where R_v is the sup-norm diameter of E_v. Good places contribute non-negatively to the Green-function version; summing the finitely many bad contributions and dividing by $nc(n)$ proves the theorem.

For $X = \mathbb{P}^1$ with the monomial basis, the determinant is the classical Vandermonde determinant and the generalized function is $\frac{1}{n(n+1)} \sum_{i \neq j} g_f(P_i, P_j)$. The special-basis construction is precisely what replaces this one-variable factorization in higher dimension.

The Lehmer-type application

The global estimate becomes quantitative arithmetic information when f extends duplication on an abelian variety.

Main theorem (Looper, Theorem 4.2)**.** *Let K be a product formula field whose completions have perfect residue fields, let A/K be an abelian variety of dimension g, and let L be symmetric and ample. There is $C = C(A, K, L) > 0$ such that every $P \in A(\overline{K}) \setminus A(\overline{K})_{\mathrm{tors}}$ which does not lie in a torsion translate of an abelian subvariety, defined after a finite extension and having everywhere potential good reduction, satisfies*

$$\widehat{h}_L(P) \geq \frac{C}{D^{\,2g+3}(\log D)^{2g}}, \qquad D = [K(P) : K] \geq 2.$$

The exclusion is essential to the method: it ensures, after reduction to the relevant geometrically simple factor, a genuinely bad non-archimedean place. In particular, a geometrically simple A with stable bad reduction somewhere satisfies the bound for every non-torsion point.

How the height bound is forced

There are four linked ingredients.

(1) For duplication, after the lift normalization and finite extension used in the proof, Corollary 3.4 makes the special transfinite diameters tend to 1. Torsion configurations with nonzero evaluation determinant are asymptotically Fekete; pluripotential equidistribution identifies their limit with the canonical measure.

(2) At a bad place v_0, that canonical measure is Haar measure on the positive-dimensional Berkovich skeleton. The tube U retracting to the identity point of the skeleton cannot contain asymptotically Fekete configurations. Along a subsequence n_ℓ, its transfinite diameter is therefore strictly smaller, giving a uniform positive local Green lower bound for $c(n_\ell)$-tuples in U.

(3) Lemma 4.1 is an interpolation statement. For a non-torsion point on a geometrically simple g-fold, among $P, 2P, \ldots, (2n^g + c(n))P$ one can choose $c(n)$ points whose evaluation determinant is nonzero. Repeated intersections of a degree-n divisor with its translates prove this without assuming genericity of the progression.

(4) Component-group and ramification bounds put suitable multiples $ke_w\Delta P$ inside U at every place over v_0, where e_w is the ramification index and Δ is the order of the relevant Néron component group. The product formula identifies the global sum of their Green functions with the average canonical height. Comparing the positive v_0-term with the universal negative error forces $n_\ell \leq BD \log D$. Since $c(n) = O(n^g)$, the largest needed multiple is $O(D^g(\log D)^g)$; quadraticity of height and $e_w \leq D$ yield the exponent $D^{2g+3}(\log D)^{2g}$.

The hardest step is the strict local diameter gap: it converts the positive-dimensional skeleton at bad reduction into a quantitative height obstruction. The theorem records a non-explicit constant, and the number-field exponent is weaker than Masser's cited bound. The paper also proves equidistribution of general asymptotically Fekete tuples (Theorem 3.5); here only the form needed for the height argument has been displayed.