Exceptional Fano varieties with small minimal log discrepancy

Louis Esser, Jihao Liu, Chengxi Wang

arXiv:2406.03570

http://arxiv.org/abs/2406.03570v1

How singular may an exceptional Fano variety be?

Work over $\mathbb{C}$. For a normal pair, the minimal log discrepancy is the infimum of log discrepancies of divisorial valuations; a variety is klt exactly when this number is positive. A klt Fano variety X is *exceptional* if (X, D) is klt for every effective $D \sim_{\mathbb{Q}} -K_X$. Equivalently here, its alpha invariant

$$\alpha(X) = \inf_{D \sim_{\mathbb{Q}} -K_X} \operatorname{lct}(X, D)$$

is greater than 1.

Let $s_0, s_1, \ldots = 2, 3, 7, 43, 1807, \ldots$ be Sylvester's sequence, $s_{n+1} = s_0 \cdots s_n + 1$. Its double-exponential growth is the source of the extreme singularities below.

Main theorem (Esser–Liu–Wang, Theorem 1.1)**.** *For every $n \geq 2$ there is an exceptional Fano n-fold X_n such that*

$$\operatorname{mld}(X_n) = \begin{cases} \dfrac{4(s_n - 1)}{s_n^3 - 9s_n}, & n \text{ even}, \\ \dfrac{4(s_n - 3)}{s_n^3 - 19s_n + 14}, & n \text{ odd}. \end{cases}$$

Hence $\operatorname{mld}(X_n) \sim 4/s_n^2$*, which tends to zero doubly exponentially with the dimension.*

The varieties are explicit well-formed, non-quasismooth hypersurfaces $X = X_d \subset \mathbb{P}(a_0, \ldots, a_{n+1})$ (Theorems 4.1–4.2), where a_i are the weights and d is the weighted degree; moreover, $-K_X = \mathcal{O}_X(1)$. The parity cases use different last weights and mixed monomials, but share the Fermat chain $x_0^2 + x_1^3 + \cdots + x_{n-1}^{s_{n-1}}$.

The surface illustrates the construction without hiding any weights:

$$X_{282} = \{x_0^2 + x_1^3 + x_2^{19}x_3 + x_1x_2x_3^5 = 0\} \subset \mathbb{P}(141, 94, 13, 35).$$

It has mld 3/35, the proven optimum among exceptional del Pezzo surfaces. In higher dimension the corresponding optimality statement is explicitly Conjecture 1.2, not part of the theorem.

Turning a Newton polytope into discrepancies

The new computational tool is Theorem 3.3. Near the unique non-quasismooth coordinate point, X_n is a cyclic quotient by $G = \mu_r$, a cyclic subgroup of the ambient torus acting freely in codimension one, of a normal Newton-nondegenerate hypersurface $Y = \{f = 0\} \subset \mathbb{A}^{n+1}$. If the quotient is well-formed and N is its toric lattice in the positive orthant $|\Delta|$, then

$$\operatorname{mld}(Y/G) = \inf_{\beta \in (N \cap |\Delta|) \setminus \{0\}} \big(\beta(x_0 \cdots x_n) - \beta(f)\big).$$

A fan refining the Newton dual fan resolves the hypersurface. The ray β extracts a toric divisor whose ambient canonical discrepancy is $\beta(x_0 \cdots x_n) - 1$, while the total transform of Y has multiplicity $\beta(f)$. Adjunction leaves exactly the difference above. Newton nondegeneracy ensures that these toroidal divisors see all discrepancies needed on a log resolution.

Why the displayed value is the global minimum

The Sylvester identities first make every defining monomial have the same weighted degree. Gcd calculations then prove the ambient space and hypersurface well-formed; the base-locus analysis gives one and only one non-quasismooth coordinate point. Normality and adjunction yield $K_X = \mathcal{O}_X(-1)$, so X is Fano once klt is known.

In the last affine chart the cyclic quotient lattice consists of integer points together with fractional points $\beta_j^i = \{j a_i / a_{n+1}\}$. Theorem 5.1 performs the parity-dependent lattice minimization in the Newton formula. It finds

$$\operatorname{mld}_x(X) = \begin{cases} (s_n - 1)/(2a_{n+1}), & n \text{ even}, \\ (s_n - 3)/(4a_{n+1}), & n \text{ odd}, \end{cases}$$

at the non-quasismooth point x. Congruences coming from weighted homogeneity identify a candidate fractional lattice point, while a case analysis shows that every other β_j gives a larger difference. This local computation alone is insufficient: Lemma 5.2 checks every remaining toric stratum and proves $\operatorname{mld}(X \setminus \{x\}) \geq 2/a_n$, which is strictly larger. Thus the candidate is the global mld. Substitution of the explicit a_{n+1} gives the two formulas in Theorem 1.1.

Why the hypersurfaces are exceptional

It remains to prove more than klt singularities. For every effective $D \sim_{\mathbb{Q}} -K_X$, the degree and the weighted-projective multiplicity bound give

$$\operatorname{mult}_p(D) \leq \frac{d}{a_n a_{n+1}}$$

at smooth points of the stack. At the exceptional affine chart, a weighted blow-up replaces X by a weighted tangent cone. Its Newton polytope contains $(1, \ldots, 1)$ in its interior, so the cone is klt; further multiplicity estimates control the pair with the transform of D. Theorems 6.5–6.6 conclude in both parity cases that

$$\alpha(X) \geq \frac{a_n a_{n+1}}{d} > 1.$$

This is precisely what makes X exceptional.

The hardest work is therefore split between two global exclusions: the lattice estimates must rule out a smaller discrepancy at every other valuation, and the alpha estimate must handle every anti-canonical divisor, especially at the non-quasismooth point. The paper gives only bounds for $\alpha(X)$, not its exact value, and claims the smallest *known* all-dimensional mlds; only the two-dimensional minimum is proved optimal.