Existence of arithmetic degrees for generic orbits and dynamical Lang-Siegel problem

Yohsuke Matsuzawa

arXiv:2407.03097

http://arxiv.org/abs/2407.03097v2

Generic orbits and arithmetic growth

Let $f\colon X \dashrightarrow X$ be a dominant rational self-map of a projective variety over $\overline{\mathbb{Q}}$. Write X_f for the points whose forward orbits avoid every indeterminacy locus, let H be ample, and set

$$\underline{\alpha}_f(x) = \liminf_n \max\{1, h_H(f^n(x))\}^{1/n}, \qquad \overline{\alpha}_f(x) = \limsup_n \max\{1, h_H(f^n(x))\}^{1/n}.$$

The orbit $O_f(x)$ is *generic* if it is infinite and meets every proper closed subset only finitely often. If $d_i(f)$ is the i-th dynamical degree, put $\mu_i(f) = d_i(f)/d_{i-1}(f)$, and set $\mu_{\dim X+1}(f) = 0$. Log-concavity gives $\mu_1 \geq \mu_2 \geq \cdots$, with $\mu_1 = d_1$.

Main theorem (Matsuzawa, Theorem 2.2)**.** *If $x \in X_f(\overline{\mathbb{Q}})$ has generic orbit, then $\alpha_f(x) = \lim_n \max\{1, h_H(f^n(x))\}^{1/n}$ exists. Moreover,*

$$\alpha_f(x) \in \{\mu_1(f), \ldots, \mu_{\dim X}(f)\}.$$

Only ratios at least 1 can occur.

This proves existence, not the full Kawaguchi–Silverman prediction $\alpha_f(x) = d_1(f)$ for a Zariski-dense orbit. Genericity is stronger than density; dynamical Mordell–Lang would make a dense orbit generic. As a consequence already available unconditionally, the paper obtains existence for every orbit of an étale endomorphism of a quasi-projective variety (Corollary 1.5).

The decisive recurrence

Choose the least ℓ with $\mu_{\ell+1} < \overline{\alpha}_f(x)$, so $\mu_{\ell+1} < \overline{\alpha}_f(x) \leq \mu_\ell$. The input [23, Theorem 3.6] is that, on a common birational model, for each $0 < \varepsilon < 1$ and every sufficiently large $m = m(\varepsilon)$, the divisor represented by

$$(f^{2m})^*H + \mu_\ell^m \mu_{\ell+1}^m H - \varepsilon^m \mu_\ell^m (f^m)^*H$$

is big. Its height is bounded below away from a proper closed set. Genericity has exactly the needed role: after discarding an initial orbit segment, all iterates avoid both that set and the indeterminacy of the chosen model.

Writing $g = f^m$, the resulting second-order height inequality is iterated in Lemma 2.4. There is $C > 0$ such that, for every $0 < \eta < 1$, after suitable shifts s, n, it gives, uniformly for $k \geq 0$,

$$h_H(g^{n+k}(f^s(x))) \geq C(\eta\mu_\ell)^{mk} h_H(g^n(f^s(x))).$$

Thus $\underline{\alpha}_f(x) \geq \eta\mu_\ell$; letting $\eta \to 1$ closes the chain $\mu_\ell \leq \underline{\alpha}_f(x) \leq \overline{\alpha}_f(x) \leq \mu_\ell$.

The product map on $X = (\mathbb{P}^1)^r$, with coordinate degrees $a_1 \geq \cdots \geq a_r > 1$, is a useful model: $\mu_i = a_i$, and a point with only its i-th coordinate non-preperiodic has arithmetic degree a_i. Such an orbit is not generic in X when $i > 1$; the example explains both the spectral ratios in the proof and why the stronger generic hypothesis matters.

From global growth to local avoidance

For a zero-dimensional subscheme Y in the paper's backward finite locus X_f^{back}, let $e(f;Y)$ be the exponential growth rate of the largest multiplicity over all inverse images of Y. The backward locus is needed because a rational map need not be finite where its inverse orbit is taken.

Main theorem (Matsuzawa, Theorem 3.3 = Theorem 1.8)**.** *Let X/K be geometrically integral and projective, let f be dominant rational, and let $x \in X_f(K)$ have generic orbit. If $Y \subset X_f^{\mathrm{back}}$ is zero-dimensional and $e(f;Y) < \alpha_f(x)$, then for every finite $S \subset M_K$,*

$$\lim_{n\to\infty} \frac{\sum_{v\in S} \lambda_{Y,v}(f^n(x))}{h_H(f^n(x))} = 0.$$

After reducing to one rational point and one place, choose an iterate $g = f^m$ so that the recurrence grows at a rate strictly larger than the backward multiplicities. Pulling the local height back through g^k replaces proximity to Y by proximity to $g^{-k}Y$. Roth's theorem bounds the latter by three times its maximal multiplicity times a global height, outside finitely many points. The multiplicity-to-growth ratio can be made arbitrarily small; the height recurrence then gives the displayed limit on every residue class modulo m.

Positive-dimensional targets and scope

For a surjective morphism, Definition 4.2 extends the notation to any closed Y by $e(f;Y) = \lim_n \max_{z\in f^{-n}(Y)} e_{f^n}(z)^{1/n}$. Theorem 4.3 (= Theorem 1.9) treats a proper closed subscheme Y for a surjective morphism of a smooth projective geometrically irreducible variety. If Y contains no positive-dimensional periodic subvariety and $e(f;Y) < \alpha_f(x)$, then any infinite set of times along which the normalized S-adic proximity has positive liminf has Banach density zero. The proof is an induction on $\dim Y$: positive Banach density supplies a shift k occurring often, and simultaneous proximity at times $n, n+k$ forces proximity to $Y \cap f^{-k}Y$, whose dimension drops because Y is not periodic.

The hypotheses mark real boundaries of the method. The first theorem needs a zero-dimensional target to invoke Roth without Vojta's conjecture; the second weakens convergence to a density-zero statement. Genericity is used to escape fixed exceptional loci, and the strict inequality $e(f;Y) < \alpha_f(x)$ is the gap that lets global height growth dominate backward ramification.